

Cheeger constants and L^2 -Betti numbers

Lewis Bowen

Workshop: Dynamics of groups and rational maps, IPAM
April 2013

The Cheeger constant

The Cheeger constant of a manifold X is defined by

$$h(X) = \inf \left\{ \frac{\text{area}(\partial M)}{\text{vol}(M)} : M \text{ compact} \subset X, \text{vol}(M) \leq \text{vol}(X)/2 \right\}.$$

The Cheeger constant

The **Cheeger constant** of a manifold X is defined by

$$h(X) = \inf \left\{ \frac{\text{area}(\partial M)}{\text{vol}(M)} : M \text{ compact } \subset X, \text{vol}(M) \leq \text{vol}(X)/2 \right\}.$$

Given a collection \mathcal{C} of subgroups of $\text{Isom}(X)$ each of which acts freely and properly discontinuously, let

$$h(X|\mathcal{C}) = \inf_{\Gamma \in \mathcal{C}} h(X/\Gamma).$$

The Cheeger constant

The **Cheeger constant** of a manifold X is defined by

$$h(X) = \inf \left\{ \frac{\text{area}(\partial M)}{\text{vol}(M)} : M \text{ compact } \subset X, \text{vol}(M) \leq \text{vol}(X)/2 \right\}.$$

Given a collection \mathcal{C} of subgroups of $\text{Isom}(X)$ each of which acts freely and properly discontinuously, let

$$h(X|\mathcal{C}) = \inf_{\Gamma \in \mathcal{C}} h(X/\Gamma).$$

Main Problem: Determine $h(X|\mathcal{C})$ in some interesting cases.

Real hyperbolic manifolds with free fundamental group

Example: What is $h(\mathbb{H}^n | \text{Free})$?

Real hyperbolic manifolds with free fundamental group

Example: What is $h(\mathbb{H}^n | \text{Free})$?

- $h(\mathbb{H}^2 | \text{Free}) = 0$.

Real hyperbolic manifolds with free fundamental group

Example: What is $h(\mathbb{H}^n | \text{Free})$?

- $h(\mathbb{H}^2 | \text{Free}) = 0.$

- $h(\mathbb{H}^3 | \text{Free}) = 0.$

Real hyperbolic manifolds with free fundamental group

Example: What is $h(\mathbb{H}^n|\text{Free})$?

- $h(\mathbb{H}^2|\text{Free}) = 0$.
- $h(\mathbb{H}^3|\text{Free}) = 0$.
- $h(\mathbb{H}^4|\text{Free}) = 0?$

Motivation: the bass note of Schottky groups

A **Schottky group** is a subgroup $\Gamma < \text{Isom}(\mathbb{H}^n)$ freely generated by elements g_1, \dots, g_m such that there are conformally round pairwise disjoint balls $B_1, \dots, B_m, B'_1, \dots, B'_m \subset \partial\mathbb{H}^n$ such that $g_i B_i = \partial\mathbb{H}^n \setminus B'_i$ for each i .

Motivation: the bass note of Schottky groups

A **Schottky group** is a subgroup $\Gamma < \text{Isom}(\mathbb{H}^n)$ freely generated by elements g_1, \dots, g_m such that there are conformally round pairwise disjoint balls $B_1, \dots, B_m, B'_1, \dots, B'_m \subset \partial\mathbb{H}^n$ such that $g_i B_i = \partial\mathbb{H}^n \setminus B'_i$ for each i .

Theorem (Philips-Sarnak, Doyle)

For every $n \geq 3$ there is a $c_n > 0$ such that if $\Gamma < \text{Isom}(\mathbb{H}^n)$ is a finitely generated Schottky group then $\lambda_0(\mathbb{H}^n/\Gamma) > c_n$.

Motivation: the bass note of Schottky groups

A **Schottky group** is a subgroup $\Gamma < \text{Isom}(\mathbb{H}^n)$ freely generated by elements g_1, \dots, g_m such that there are conformally round pairwise disjoint balls $B_1, \dots, B_m, B'_1, \dots, B'_m \subset \partial\mathbb{H}^n$ such that $g_i B_i = \partial\mathbb{H}^n \setminus B'_i$ for each i .

Theorem (Philips-Sarnak, Doyle)

For every $n \geq 3$ there is a $c_n > 0$ such that if $\Gamma < \text{Isom}(\mathbb{H}^n)$ is a finitely generated Schottky group then $\lambda_0(\mathbb{H}^n/\Gamma) > c_n$.

Cheeger: $\lambda_0(X) \geq h(X)^2/4$.

Motivation: the bass note of Schottky groups

A **Schottky group** is a subgroup $\Gamma < \text{Isom}(\mathbb{H}^n)$ freely generated by elements g_1, \dots, g_m such that there are conformally round pairwise disjoint balls $B_1, \dots, B_m, B'_1, \dots, B'_m \subset \partial\mathbb{H}^n$ such that $g_i B_i = \partial\mathbb{H}^n \setminus B'_i$ for each i .

Theorem (Philips-Sarnak, Doyle)

For every $n \geq 3$ there is a $c_n > 0$ such that if $\Gamma < \text{Isom}(\mathbb{H}^n)$ is a finitely generated Schottky group then $\lambda_0(\mathbb{H}^n/\Gamma) > c_n$.

Cheeger: $\lambda_0(X) \geq h(X)^2/4$.

Buser: $\lambda_0(X) \leq c_1 h(X) + c_2 h(X)^2$ where $c_1, c_2 > 0$ depend on dimension and a lower bound on Ricci curvature.

Motivation: the bass note of Schottky groups

A **Schottky group** is a subgroup $\Gamma < \text{Isom}(\mathbb{H}^n)$ freely generated by elements g_1, \dots, g_m such that there are conformally round pairwise disjoint balls $B_1, \dots, B_m, B'_1, \dots, B'_m \subset \partial\mathbb{H}^n$ such that $g_i B_i = \partial\mathbb{H}^n \setminus B'_i$ for each i .

Theorem (Philips-Sarnak, Doyle)

For every $n \geq 3$ there is a $c_n > 0$ such that if $\Gamma < \text{Isom}(\mathbb{H}^n)$ is a finitely generated Schottky group then $\lambda_0(\mathbb{H}^n/\Gamma) > c_n$.

Cheeger: $\lambda_0(X) \geq h(X)^2/4$.

Buser: $\lambda_0(X) \leq c_1 h(X) + c_2 h(X)^2$ where $c_1, c_2 > 0$ depend on dimension and a lower bound on Ricci curvature.

$\Rightarrow h(\mathbb{H}^n | \text{Schottky}) > 0$ for $n \geq 3$.

Motivation: fiber bundles

Problem: does there exist a finite-volume real hyperbolic 4-manifold which fibers over a surface with a fiber a surface?

Motivation: fiber bundles

Problem: does there exist a finite-volume real hyperbolic 4-manifold which fibers over a surface with a fiber a surface?

[U. Hamenstädt : not if the fiber and the base are closed surfaces.]

Motivation: fiber bundles

Problem: does there exist a finite-volume real hyperbolic 4-manifold which fibers over a surface with a fiber a surface?

[U. Hamenstädt : not if the fiber and the base are closed surfaces.]

If so, the fundamental group of the fiber surface corresponds to a subgroup $\Gamma < Isom(X)$ with a limit set which is all of ∂X .

Motivation: fiber bundles

Problem: does there exist a finite-volume real hyperbolic 4-manifold which fibers over a surface with a fiber a surface?

[U. Hamenstädt : not if the fiber and the base are closed surfaces.]

If so, the fundamental group of the fiber surface corresponds to a subgroup $\Gamma < Isom(X)$ with a limit set which is all of ∂X .

Is it possible to deform such a subgroup so that it is geometrically finite and has limit set with Hausdorff dimension close to 3?

Motivation: fiber bundles

Problem: does there exist a finite-volume real hyperbolic 4-manifold which fibers over a surface with a fiber a surface?

[U. Hamenstädt : not if the fiber and the base are closed surfaces.]

If so, the fundamental group of the fiber surface corresponds to a subgroup $\Gamma < Isom(X)$ with a limit set which is all of ∂X .

Is it possible to deform such a subgroup so that it is geometrically finite and has limit set with Hausdorff dimension close to 3?

If so then $h(\mathbb{H}^4 | \text{Surface}) = 0$.

(PS theory $\Rightarrow \lambda_0(\mathbb{H}^n / \Gamma) = \dim(L\Gamma)(n - 1 - \dim(L\Gamma))$)

Main results

Definition

Let \mathcal{G}_d be the class of residually finite groups Γ with

$$b_d^{(2)}(\Gamma') = 0 \quad \forall \Gamma' < \Gamma.$$

Main results

Definition

Let \mathcal{G}_d be the class of residually finite groups Γ with
 $b_d^{(2)}(\Gamma') = 0 \forall \Gamma' < \Gamma$.

Theorem

Let X be a contractible smooth complete Riemannian manifold. Suppose there is a residually finite cocompact lattice $\Lambda < \text{Isom}(X)$ such that $b_d^{(2)}(\Lambda) > 0$. Then $h(X|\mathcal{G}_d) > 0$.

Main results

Definition

Let \mathcal{G}_d be the class of residually finite groups Γ with $b_d^{(2)}(\Gamma') = 0 \forall \Gamma' < \Gamma$.

Theorem

Let X be a contractible smooth complete Riemannian manifold. Suppose there is a residually finite cocompact lattice $\Lambda < \text{Isom}(X)$ such that $b_d^{(2)}(\Lambda) > 0$. Then $h(X|\mathcal{G}_d) > 0$.

Corollary

For any even number $n \geq 2$, $h(\mathbb{H}^n|\mathcal{G}_{n/2}) > 0$.

Main results

Definition

Let \mathcal{G}_d be the class of residually finite groups Γ with $b_d^{(2)}(\Gamma') = 0 \forall \Gamma' < \Gamma$.

Theorem

Let X be a contractible smooth complete Riemannian manifold. Suppose there is a residually finite cocompact lattice $\Lambda < \text{Isom}(X)$ such that $b_d^{(2)}(\Lambda) > 0$. Then $h(X|\mathcal{G}_d) > 0$.

Corollary

For any even number $n \geq 2$, $h(\mathbb{H}^n|\mathcal{G}_{n/2}) > 0$.

Lemma

If Γ is isomorphic to a subgroup of a fundamental group of a complete finite volume hyperbolic 3-manifold then $\Gamma \in \mathcal{G}_d$ for all $d \geq 2$. So $h(\mathbb{H}^{2n}|\text{Free}) > 0$ for all $n \geq 2$.

Benjamini-Schramm convergence

Let $\{K_j\}$ be a sequence of finite connected simplicial complexes.

Benjamini-Schramm convergence

Let $\{K_j\}$ be a sequence of finite connected simplicial complexes. Given a finite connected simplicial complex L and a vertex v in L , let

$$\rho_j(L, v) = \#\{w \in V(K_j) : \exists \text{ embedding } \phi : L \rightarrow K_j \text{ with } \phi(v) = w\}.$$

Benjamini-Schramm convergence

Let $\{K_i\}$ be a sequence of finite connected simplicial complexes. Given a finite connected simplicial complex L and a vertex v in L , let

$$\rho_i(L, v) = \#\{w \in V(K_i) : \exists \text{ embedding } \phi : L \rightarrow K_i \text{ with } \phi(v) = w\}.$$

We say $\{K_i\}$ is *Benjamini-Schramm convergent* if

$$\lim_{i \rightarrow \infty} \frac{\rho_i(L, v)}{|V(K_i)|} =: \bar{\rho}(L, v)$$

exists for every (L, v) .

Benjamini-Schramm convergence

Let $\{K_i\}$ be a sequence of finite connected simplicial complexes. Given a finite connected simplicial complex L and a vertex v in L , let

$$\rho_i(L, v) = \#\{w \in V(K_i) : \exists \text{ embedding } \phi : L \rightarrow K_i \text{ with } \phi(v) = w\}.$$

We say $\{K_i\}$ is *Benjamini-Schramm convergent* if

$$\lim_{i \rightarrow \infty} \frac{\rho_i(L, v)}{|V(K_i)|} =: \bar{\rho}(L, v)$$

exists for every (L, v) .

Lemma (G. Elek, 2010)

Suppose $\{K_i\}$ is BS-convergent and suppose there is a uniform bound on the vertex degrees of $\{K_i\}$. Then for every $d > 0$,

$$\lim_{i \rightarrow \infty} \frac{b_d(K_i)}{|V(K_i)|}$$

exists.

Lück Approximation

Theorem (W. Lück, 1994)

Let Λ be a finitely generated group, $\{\Lambda_n\}_{n=1}^{\infty}$ a decreasing sequence of finite-index normal subgroups of Λ with $\bigcap_n \Lambda_n = \{e\}$. Then

$$\lim_{n \rightarrow \infty} \frac{b_d(\Lambda_n)}{[\Lambda : \Lambda_n]} = b_d^{(2)}(\Lambda).$$

Another view of Benjamini-Schramm convergence

A *rooted simplicial complex* is a pair (K, v) where K is a connected simplicial complex and $v \in V(K)$.

Another view of Benjamini-Schramm convergence

A *rooted simplicial complex* is a pair (K, v) where K is a connected simplicial complex and $v \in V(K)$.

Let $\text{RSC}(\Delta)$ be the space of all root-isomorphism classes of rooted simplicial complex with vertex degree bound Δ . There's a natural topology on $\text{RSC}(\Delta)$.

Another view of Benjamini-Schramm convergence

A *rooted simplicial complex* is a pair (K, v) where K is a connected simplicial complex and $v \in V(K)$.

Let $\text{RSC}(\Delta)$ be the space of all root-isomorphism classes of rooted simplicial complex with vertex degree bound Δ . There's a natural topology on $\text{RSC}(\Delta)$.

Given a finite connected simplicial complex K , define

$$\mu_K = \sum_{v \in V(K)} \delta_{[K, v]}.$$

Lemma

$\{K_i\}$ is BS-convergent $\Leftrightarrow \mu_{K_i}$ converges in the weak* topology on $\mathcal{M}_1(\text{RSC}(\Delta))$.

Metric measure spaces

A *metric measure space* is a (proper separable complete) metric space with a (Radon) measure: $M = (M, \text{dist}_M, \text{vol}_M)$.

Metric measure spaces

A *metric measure space* is a (proper separable complete) metric space with a (Radon) measure: $M = (M, \text{dist}_M, \text{vol}_M)$.

A *pointed mm-space* is a pair (M, ρ) where M is an mm-space and $\rho \in M$.

Metric measure spaces

A *metric measure space* is a (proper separable complete) metric space with a (Radon) measure: $M = (M, \text{dist}_M, \text{vol}_M)$.

A *pointed mm-space* is a pair (M, ρ) where M is an mm-space and $\rho \in M$.

Let \mathbb{M} denote the set of isomorphism classes of pointed mm-spaces.

Metric measure spaces

A *metric measure space* is a (proper separable complete) metric space with a (Radon) measure: $M = (M, \text{dist}_M, \text{vol}_M)$.

A *pointed mm-space* is a pair (M, p) where M is an mm-space and $p \in M$.

Let \mathbb{M} denote the set of isomorphism classes of pointed mm-spaces.

A sequence $[M_i, p_i] \in \mathbb{M}$ converges to $[M_\infty, p_\infty]$ iff there exist a complete separable metric space Z and isometric embeddings $\varphi_i : M_i \rightarrow Z$ ($1 \leq i \leq \infty$) such that

- $(\varphi_i(M_i), \varphi_i(p_i))$ converges to $(\varphi_\infty(M_\infty), \varphi_\infty(p_\infty))$ in the pointed Hausdorff topology;
- $(\varphi_i)_* \text{vol}_{M_i}$ converges to $(\varphi_\infty)_* \text{vol}_{M_\infty}$ in the weak* topology.

A variation on Elek's Theorem

Let $\mathcal{M}_1(\mathbb{M})$ = the space of Borel probability measures on \mathbb{M} .

Theorem

Let M_i be a sequence of path-connected finite-volume mm-spaces. Let μ_i be the push-forward of vol_{M_i} under the map $p \mapsto [M_i, p]$. Suppose μ_i converges in $\mathcal{M}_1(\mathbb{M})$ (weak*). Suppose also that there exist constants $\epsilon, v_0, v_1, \Delta > 0$ such that

-
-
-
-
-

Then

$$\lim_{i \rightarrow \infty} \frac{b_d(M_i)}{\text{vol}_{M_i}(M_i)} \text{ exists for every } d > 0.$$

A variation on Elek's Theorem

Let $\mathcal{M}_1(\mathbb{M})$ = the space of Borel probability measures on \mathbb{M} .

Theorem

Let M_i be a sequence of path-connected finite-volume mm-spaces. Let μ_i be the push-forward of vol_{M_i} under the map $p \mapsto [M_i, p]$. Suppose μ_i converges in $\mathcal{M}_1(\mathbb{M})$ (weak*). Suppose also that there exist constants $\epsilon, \nu_0, \nu_1, \Delta > 0$ such that

- vol_{M_i} is purely non-atomic and fully supported;
- every sphere in M_i has measure zero;
- $0 < \nu_0 \leq \text{vol}_{M_i}(B_{M_i}(p, \epsilon/2)) \leq \text{vol}_{M_i}(B_{M_i}(p, 10\epsilon)) \leq \nu_1 \forall p \in M_i$;
- $B_{M_i}(p, r)$ is strongly convex for every $r \leq 10\epsilon$;
- the max. cardinality of an ϵ -separated subset of $B_{M_i}(p, 20\epsilon)$ is $\leq \Delta$.

Then

$$\lim_{i \rightarrow \infty} \frac{b_d(M_i)}{\text{vol}_{M_i}(M_i)} \text{ exists for every } d > 0.$$

Proof Sketch of the Main Theorem

Definition

Let \mathcal{G}_d be the class of residually finite groups Γ with
 $b_d^{(2)}(\Gamma') = 0 \forall \Gamma' < \Gamma$.

Proof Sketch of the Main Theorem

Definition

Let \mathcal{G}_d be the class of residually finite groups Γ with $b_d^{(2)}(\Gamma') = 0 \forall \Gamma' < \Gamma$.

Theorem

Let X be a contractible smooth complete Riemannian manifold. Suppose there is a residually finite cocompact lattice $\Lambda < \text{Isom}(X)$ such that $b_d^{(2)}(\Lambda) > 0$. Then $h(X|\mathcal{G}_d) > 0$.

Proof Sketch of the Main Theorem

Definition

Let \mathcal{G}_d be the class of residually finite groups Γ with $b_d^{(2)}(\Gamma') = 0 \forall \Gamma' < \Gamma$.

Theorem

Let X be a contractible smooth complete Riemannian manifold. Suppose there is a residually finite cocompact lattice $\Lambda < \text{Isom}(X)$ such that $b_d^{(2)}(\Lambda) > 0$. Then $h(X|\mathcal{G}_d) > 0$.

It suffices to show that if $\Gamma_i < \text{Isom}(X)$ is a sequence of residually finite subgroups which act freely and properly discontinuous and $h(X/\Gamma_i) \rightarrow 0$ then there exist subgroups $\Gamma'_i < \Gamma_i$ s.t. $b_d^{(2)}(\Gamma'_i) > 0$ for all but finitely many i .

A very rough idea

Let $M_i \subset X/\Gamma_i$ be a compact connected submanifold with

$$\lim_{i \rightarrow \infty} \frac{\text{area}(\partial M_i)}{\text{vol}(M_i)} = 0.$$

A very rough idea

Let $M_i \subset X/\Gamma_i$ be a compact connected submanifold with

$$\lim_{i \rightarrow \infty} \frac{\text{area}(\partial M_i)}{\text{vol}(M_i)} = 0.$$

Let us pretend that $\{M_i\}$ Benjamini-Schramm converges to X . Then

$$\lim_{i \rightarrow \infty} \frac{b_d(M_i)}{\text{vol}(M_i)} = \frac{b_d^{(2)}(\Lambda)}{\text{vol}(X/\Lambda)} > 0.$$

A very rough idea

Let $M_i \subset X/\Gamma_i$ be a compact connected submanifold with

$$\lim_{i \rightarrow \infty} \frac{\text{area}(\partial M_i)}{\text{vol}(M_i)} = 0.$$

Let us pretend that $\{M_i\}$ Benjamini-Schramm converges to X . Then

$$\lim_{i \rightarrow \infty} \frac{b_d(M_i)}{\text{vol}(M_i)} = \frac{b_d^{(2)}(\Lambda)}{\text{vol}(X/\Lambda)} > 0.$$

Let us also pretend that $\pi_1(M_i) \rightarrow \Gamma_i$ is injective. Then $b_d(M_i) = b_d(\Gamma'_i)$ for some $\Gamma'_i < \Gamma_i$.

A very rough idea

Let $M_i \subset X/\Gamma_i$ be a compact connected submanifold with

$$\lim_{i \rightarrow \infty} \frac{\text{area}(\partial M_i)}{\text{vol}(M_i)} = 0.$$

Let us pretend that $\{M_i\}$ Benjamini-Schramm converges to X . Then

$$\lim_{i \rightarrow \infty} \frac{b_d(M_i)}{\text{vol}(M_i)} = \frac{b_d^{(2)}(\Lambda)}{\text{vol}(X/\Lambda)} > 0.$$

Let us also pretend that $\pi_1(M_i) \rightarrow \Gamma_i$ is injective. Then $b_d(M_i) = b_d(\Gamma'_i)$ for some $\Gamma'_i < \Gamma_i$.

By Lück Approximation, we may pass to a finite-index subgroup $\Gamma''_i < \Gamma'_i$ such that

$$\frac{b_d(\Gamma''_i)}{[\Gamma'_i : \Gamma''_i]} \approx b_d^{(2)}(\Gamma'_i).$$

It follows that $b_d^{(2)}(\Gamma'_i) > 0$ for all but finitely many i .

Let $M_i \subset X/\Gamma_i$ be a compact connected submanifold with

$$\lim_{i \rightarrow \infty} \frac{\text{area}(\partial M_i)}{\text{vol}(M_i)} = 0.$$

Let $M_i \subset X/\Gamma_i$ be a compact connected submanifold with

$$\lim_{i \rightarrow \infty} \frac{\text{area}(\partial M_i)}{\text{vol}(M_i)} = 0.$$

Buser's Hair-cutting Lemma \implies we can choose M_i so that

$$\lim_{i \rightarrow \infty} \frac{\text{vol}(N_r(\partial M_i))}{\text{vol}(M_i)} = 0 \quad \forall r > 0.$$

Let $M_i \subset X/\Gamma_i$ be a compact connected submanifold with

$$\lim_{i \rightarrow \infty} \frac{\text{area}(\partial M_i)}{\text{vol}(M_i)} = 0.$$

Buser's Hair-cutting Lemma \implies we can choose M_i so that

$$\lim_{i \rightarrow \infty} \frac{\text{vol}(N_r(\partial M_i))}{\text{vol}(M_i)} = 0 \quad \forall r > 0.$$

Let $\Gamma'_i < \Gamma_i$ be the image of $\pi_1(M_i)$ under the inclusion $M_i \rightarrow X/\Gamma_i$.

Let $M_i \subset X/\Gamma_i$ be a compact connected submanifold with

$$\lim_{i \rightarrow \infty} \frac{\text{area}(\partial M_i)}{\text{vol}(M_i)} = 0.$$

Buser's Hair-cutting Lemma \implies we can choose M_i so that

$$\lim_{i \rightarrow \infty} \frac{\text{vol}(N_r(\partial M_i))}{\text{vol}(M_i)} = 0 \quad \forall r > 0.$$

Let $\Gamma'_i < \Gamma_i$ be the image of $\pi_1(M_i)$ under the inclusion $M_i \rightarrow X/\Gamma_i$.

Lift M_i to a subset $M'_i \subset X/\Gamma'_i$. We still have

$$\lim_{i \rightarrow \infty} \frac{\text{vol}(N_r(\partial M'_i))}{\text{vol}(M'_i)} = 0 \quad \forall r > 0.$$

- \exists finite-index subgroup $\Gamma_i'' < \Gamma_i'$ such that
- if M_i'' is the lift of M_i' to X/Γ_i'' then M_i'' has no short homotopically-nontrivial loops.

- \exists finite-index subgroup $\Gamma_i'' < \Gamma_i'$ such that
- if M_i'' is the lift of M_i' to X/Γ_i'' then M_i'' has no short homotopically-nontrivial loops.

We still have

$$\lim_{i \rightarrow \infty} \frac{\text{vol}(N_r(\partial M_i''))}{\text{vol}(M_i'')} = 0 \quad \forall r > 0.$$

So $\{M_i''\}$ Benjamini-Schramm converges to X .

- \exists finite-index subgroup $\Gamma_i'' < \Gamma_i'$ such that
- if M_i'' is the lift of M_i' to X/Γ_i'' then M_i'' has no short homotopically-nontrivial loops.

We still have

$$\lim_{i \rightarrow \infty} \frac{\text{vol}(N_r(\partial M_i''))}{\text{vol}(M_i'')} = 0 \quad \forall r > 0.$$

So $\{M_i''\}$ Benjamini-Schramm converges to X .

By [Lück Approximation](#), we can also choose Γ_i'' so that

$$\liminf_{i \rightarrow \infty} \frac{b_d(\Gamma_i'')}{\text{vol}(M_i'')} = \liminf_{i \rightarrow \infty} \frac{b_d^{(2)}(\Gamma_i')}{\text{vol}(M_i')}.$$

Benjamini-Schramm convergence implies:

$$\lim_{i \rightarrow \infty} \frac{b_d(M_i'')}{\text{vol}(M_i'')} = \frac{b_d^{(2)}(\Lambda)}{\text{vol}(X/\Lambda)} > 0.$$

If $b_d(M_i'') = b_d(\Gamma_i'')$ then we're done.

Benjamini-Schramm convergence implies:

$$\lim_{i \rightarrow \infty} \frac{b_d(M_i'')}{\text{vol}(M_i'')} = \frac{b_d^{(2)}(\Lambda)}{\text{vol}(X/\Lambda)} > 0.$$

If $b_d(M_i'') = b_d(\Gamma_i'')$ then we're done.

The [Mayer-Vietoris sequence](#)

$$\cdots \rightarrow H_d(\partial M_i'') \rightarrow H_d(M_i'') \oplus H_d(X/\Gamma_i'' - M_i'') \rightarrow H_d(X/\Gamma_i'') \rightarrow \cdots$$

implies

$$b_d(M_i'') \leq b_d(\Gamma_i'') + b_d(\partial M_i'').$$

Benjamini-Schramm convergence implies:

$$\lim_{i \rightarrow \infty} \frac{b_d(M_i'')}{\text{vol}(M_i'')} = \frac{b_d^{(2)}(\Lambda)}{\text{vol}(X/\Lambda)} > 0.$$

If $b_d(M_i'') = b_d(\Gamma_i'')$ then we're done.

The [Mayer-Vietoris sequence](#)

$$\cdots \rightarrow H_d(\partial M_i'') \rightarrow H_d(M_i'') \oplus H_d(X/\Gamma_i'' - M_i'') \rightarrow H_d(X/\Gamma_i'') \rightarrow \cdots$$

implies

$$b_d(M_i'') \leq b_d(\Gamma_i'') + b_d(\partial M_i'').$$

But $b_d(\partial M_i'') \leq C \text{vol}(N_r(\partial M_i''))$ for some $r, C > 0$.

Benjamini-Schramm convergence implies:

$$\lim_{i \rightarrow \infty} \frac{b_d(M_i'')}{\text{vol}(M_i'')} = \frac{b_d^{(2)}(\Lambda)}{\text{vol}(X/\Lambda)} > 0.$$

If $b_d(M_i'') = b_d(\Gamma_i'')$ then we're done.

The [Mayer-Vietoris sequence](#)

$$\cdots \rightarrow H_d(\partial M_i'') \rightarrow H_d(M_i'') \oplus H_d(X/\Gamma_i'' - M_i'') \rightarrow H_d(X/\Gamma_i'') \rightarrow \cdots$$

implies

$$b_d(M_i'') \leq b_d(\Gamma_i'') + b_d(\partial M_i'').$$

But $b_d(\partial M_i'') \leq C \text{vol}(N_r(\partial M_i''))$ for some $r, C > 0$. So

$$\frac{b_d(\partial M_i'')}{\text{vol}(M_i'')} \rightarrow 0.$$

Benjamini-Schramm convergence implies:

$$\liminf_{i \rightarrow \infty} \frac{b_d^{(2)}(\Gamma'_i)}{\text{vol}(M'_i)} = \liminf_{i \rightarrow \infty} \frac{b_d(\Gamma''_i)}{\text{vol}(M''_i)} \geq \lim_{i \rightarrow \infty} \frac{b_d(M''_i)}{\text{vol}(M''_i)} = \frac{b_d^{(2)}(\Lambda)}{\text{vol}(X/\Lambda)} > 0.$$

If $b_d(M''_i) = b_d(\Gamma''_i)$ then we're done.

The Mayer-Vietoris sequence

$$\cdots \rightarrow H_d(\partial M''_i) \rightarrow H_d(M''_i) \oplus H_d(X/\Gamma''_i - M''_i) \rightarrow H_d(X/\Gamma''_i) \rightarrow \cdots$$

implies

$$b_d(M''_i) \leq b_d(\Gamma''_i) + b_d(\partial M''_i).$$

But $b_d(\partial M''_i) \leq C \text{vol}(N_r(\partial M''_i))$ for some $r, C > 0$. So

$$\frac{b_d(\partial M''_i)}{\text{vol}(M''_i)} \rightarrow 0.$$

Open Questions

- Is $h(\mathbb{H}^n | \text{Free}) > 0$ when $n > 3$ is odd?

Open Questions

- Is $h(\mathbb{H}^n | \text{Free}) > 0$ when $n > 3$ is odd?
- Is $h(\mathbb{H}^4 | \text{Free})$ attained?

Open Questions

- Is $h(\mathbb{H}^n | \text{Free}) > 0$ when $n > 3$ is odd?
- Is $h(\mathbb{H}^4 | \text{Free})$ attained? by a finitely generated free group?

Open Questions

- Is $h(\mathbb{H}^n | \text{Free}) > 0$ when $n > 3$ is odd?
- Is $h(\mathbb{H}^4 | \text{Free})$ attained? by a finitely generated free group?
- If $\Gamma < \text{Isom}(\mathbb{H}^n)$ is the fundamental group of a fiber in a fiber bundle over an $(n - 2)$ -manifold, does $h(\mathbb{H}^n / \Gamma)$ realize $h(\mathbb{H}^n | \text{deformations}(\text{fund. gp. fiber}))$?

Open Questions

- Is $h(\mathbb{H}^n | \text{Free}) > 0$ when $n > 3$ is odd?
- Is $h(\mathbb{H}^4 | \text{Free})$ attained? by a finitely generated free group?
- If $\Gamma < \text{Isom}(\mathbb{H}^n)$ is the fundamental group of a fiber in a fiber bundle over an $(n - 2)$ -manifold, does $h(\mathbb{H}^n / \Gamma)$ realize $h(\mathbb{H}^n | \text{deformations}(\text{fund. gp. fiber}))$?
- Suppose $\Gamma < \text{Isom}(\mathbb{H}^4)$ is discrete, properly discontinuous with infinite covolume and $b_2^{(2)}(\Gamma) > 0$. Is $h(\mathbb{H}^4 | \text{deformations}(\Gamma)) = 0$?

Open Questions

- Is $h(\mathbb{H}^n | \text{Free}) > 0$ when $n > 3$ is odd?
- Is $h(\mathbb{H}^4 | \text{Free})$ attained? by a finitely generated free group?
- If $\Gamma < \text{Isom}(\mathbb{H}^n)$ is the fundamental group of a fiber in a fiber bundle over an $(n - 2)$ -manifold, does $h(\mathbb{H}^n / \Gamma)$ realize $h(\mathbb{H}^n | \text{deformations}(\text{fund. gp. fiber}))$?
- Suppose $\Gamma < \text{Isom}(\mathbb{H}^4)$ is discrete, properly discontinuous with infinite covolume and $b_2^{(2)}(\Gamma) > 0$. Is $h(\mathbb{H}^4 | \text{deformations}(\Gamma)) = 0$?
- What is $h(\mathbb{H}^4 | \mathcal{G}_2)$?