

Monte Carlo methods for kinetic equations

Lecture 4: Hybrid methods and variance reduction

Lorenzo Pareschi

Department of Mathematics & CMCS
University of Ferrara
Italy



<http://utenti.unife.it/lorenzo.pareschi/>
lorenzo.pareschi@unife.it

KT2009: Tutorials
IPAM, March 10-13, 2009

Outline

1 Introduction

- Multiscale problems and methods
- Kinetic equations

2 Hybrid methods

- Hybrid representation
- The hybrid method
- Macroscopic solvers
- Numerical results


3 Hydro-guided Monte Carlo

- Basic principles
- The HG method
- Numerical results

Multiscale problems

- A broad range of scientific problems involve multiple scales and multi-scale phenomena (material science, chemistry, fluid dynamics, biology...). These involve **different physical laws** which govern the processes at **different scales**.
- On the computational side, several important classes of numerical methods have been developed which address explicitly the multiscale nature of the solutions (**wavelets, multigrid, domain decomposition, stiff solvers, adaptive mesh refinements...**).
- For many problems, representation or solution on the fine-scale is impossible because of the **overwhelming costs**.

The different scales



Multiscale methods

- **Couplings** of atomistic or molecular, and more generally microscopic **stochastic models**, to macroscopic **deterministic models** based on ODEs and PDEs is highly desirable in many applications. Similar arguments apply also to numerical methods¹.
- A classical field where this coupling play an important rule is that of **kinetic equations**. In such system the time scale is proportional to a relaxation time ε and a strong model (and dimension) reduction is obtained when $\varepsilon \rightarrow 0$.
- Many **examples** could depict this situation, **rarefied gas dynamics**, **plasma physics**, **granular gases**, **turbulence**,...

¹W.E, B.Engquist CMS '03, N. AMS '03

Kinetic equations

Kinetic equations

$$\partial_t f + v \nabla_x f = \frac{1}{\varepsilon} Q(f, f), \quad x, v \in \mathbb{R}^d, d \geq 1, \quad (\text{microscale})$$

Here $f = f(x, v, t) \geq 0$ is the particle density and $Q(f, f)$ describes the particle interactions. In rarefied gas dynamics the equilibrium functions M for which $Q(M, M) = 0$ are local Maxwellian

$$M(\rho, u, T)(v) = \frac{\rho}{(2\pi T)^{d/2}} \exp\left(-\frac{|u - v|^2}{2T}\right),$$

where we define the density, mean velocity and temperature as

$$\rho = \int_{\mathbb{R}^d} f \, dv, \quad u = \frac{1}{\rho} \int_{\mathbb{R}^d} v f \, dv, \quad T = \frac{1}{d\rho} \int_{\mathbb{R}^d} [v - u]^2 f \, dv.$$

Fluid limit

If we multiply the kinetic equation by its **collision invariants** $(1, v, |v|^2)$ and integrate the result in velocity space we obtain five equations that describe the **balance of mass, momentum and energy**. The system is not closed since it involves higher order moments of the distribution function f .

As $\varepsilon \rightarrow 0$ we have $Q(f, f) \rightarrow 0$ and thus f approaches the local Maxwellian M_f . Higher order moments of f can be computed as function of ρ , u , and T and we obtain the closed system

Compressible Euler equations

$$\begin{cases} \partial_t \varrho + \nabla_x \cdot (\varrho u) = 0, \\ \partial_t \varrho u + \nabla_x \cdot (\varrho u \otimes u + p) = 0, \\ \partial_t E + \nabla_x \cdot (u(E + p)) = 0, \end{cases} \quad (\text{macroscale})$$


where p is the gas pressure.

Generalizations

- The **macroscale process** is described by the conserved quantities $U = (\rho, u, T)$ whereas the **microscale process** is described by f . The two processes and state variables are related by compression and reconstruction operators P and R , such that $P(f) = U$ and $R(U) = f$, with the property $PR = I$, where I is the identity operator.
- The **compression operator** is a projection to low order moments. The **reconstruction operator** does the opposite and it is under-determined, except close to the local equilibrium state when $Q(f, f) = 0$ implies $f = M(U)$.

Hybrid representation

The solution is represented at each space point as a combination of a **nonequilibrium part** (microscale) and an **equilibrium part** (macroscale)



The starting point is the following²

Definition II - hybrid function

Given a probability density $f(v)$, $v \in \mathbb{R}^d$ (i.e. $f(v) \geq 0$, $\int f(v)dv = 1$) and a probability density $M(v)$, $v \in \mathbb{R}^d$ called equilibrium density, we define $w(v) \in [0, 1]$ and $\tilde{f}(v) \geq 0$ in the following way

$$w(v) = \begin{cases} \frac{f(v)}{M(v)}, & f(v) \leq M(v) \neq 0 \\ 1, & f(v) \geq M(v) \end{cases}$$

and $\tilde{f}(v) = f(v) - w(v)M(v)$. Thus $f(v)$ can be represented as

$$f(v) = \tilde{f}(v) + w(v)M(v).$$

²L.P. ESAIM '05, L.P., G.Dimarco CMS '06, MMS '08

Taking $\beta = \min_v \{w(v)\}$, and $\tilde{f}(v) = f(v) - \beta M(v)$, we have

$$\int \tilde{f}(v) dv = 1 - \beta.$$

Let us define for $\beta \neq 1$ the probability density

$$f_p(v) = \frac{\tilde{f}(v)}{1 - \beta}.$$

The case $\beta = 1$ is trivial since it implies $f \equiv M$. Thus we recover the hybrid representation³ as

$$f(v) = (1 - \beta)f_p(v) + \beta M(v).$$

³ R.E.Caffisch, L.P. JCP '99

The general methodology

Now we consider the following general representation

$$f(x, v, t) = \underbrace{\tilde{f}(x, v, t)}_{\text{nonequilibrium}} + \underbrace{w(x, v, t)M(\rho(x, t), u(x, t), T(x, t))(v)}_{\text{equilibrium}}.$$

The nonequilibrium part $\tilde{f}(x, v, t)$ is represented stochastically, whereas the equilibrium part $w(x, v, t)M(\rho(x, t), u(x, t), T(x, t))(v)$ deterministically. The general methodology is the following.

- Solve the evolution of the non equilibrium part by **Monte Carlo methods**. Thus $\tilde{f}(x, v, t)$ is represented by a set of samples (particles) in the computational domain.
- Solve the evolution of the equilibrium part by **deterministic methods**. Thus $w(x, v, t)M(\rho(x, t), u(x, t), T(x, t))(v)$ is obtained from a suitable grid in the computational domain.

The general methodology

The starting point of the method is the classical operator splitting which consists in solving first a homogeneous **collision step**

$$(C) \quad \partial_t f^r(x, v, t) = \frac{1}{\varepsilon} Q(f^r, f^r)(x, v, t)$$

and then a free **transport step**

$$(T) \quad \partial_t f^c(x, v, t) + v \cdot \nabla_x f^c(x, v, t) = 0.$$

Except for BGK-like models where the collision term has the form $Q(f, f) = M - f$, one needs a suitable solver for the stiff nonlinear collision operator⁴.

⁴ E.Gabetta, L.P., G.Toscani SINUM '97

Sketch of the basic method

- C:** Starting from a hybrid function $f(t) = \tilde{f}(t) + w(t)M(t)$ solve the collision step $f^r(t + \Delta t) = \lambda f(t) + (1 - \lambda)M(t)$ with $\lambda = e^{-\Delta t/\varepsilon}$.
- 1 The new value $\tilde{f}^r(t + \Delta t) = \lambda \tilde{f}(t)$ is computed by particles.
 - 2 Set $w^r(t + \Delta t) = \lambda w(t) + 1 - \lambda$.
 - 3 Discard a fraction of Monte Carlo samples since $w^r(t + \Delta t) \geq w(t)$.
- T:** Starting from the hybrid function $f^r(t + \Delta t)$ computed above solve the transport step $f(x, v, t + \Delta t) = f^r(x - v\Delta t, v, t + \Delta t)$.
- 1 Transport the particle fraction $\tilde{f}^r(x - v\Delta t, v, t + \Delta t)$ by simple particles shifts.
 - 2 Transport the deterministic fraction $w^r(x - v\Delta t, v, t + \Delta t)M(x - v\Delta t, v, t)$ by a deterministic scheme.
 - 3 Project the computed solution to the hybrid form $f(t + \Delta t) = \tilde{f}(t + \Delta t) + w(t + \Delta t)M(t + \Delta t)$.

Remarks

Note that point 2 of the transport corresponds to a Maxwellian shift analogous to that usually performed in the so called kinetic or Boltzmann schemes for the Euler equations⁵.

Clearly point 3 after the transport step is crucial for the details of the hybrid method. We have considered three different possible reconstructions

(0) We loose entirely equilibrium thus $w(x, v, t + \Delta t) = 0$.

(C) We compute the new equilibrium fraction from $w^r(x - v\Delta t, v, t + \Delta t)M(x - v\Delta t, v, t)$ using definition I.

(1) We compute the new equilibrium fraction from $w^r(x - v\Delta t, v, t + \Delta t)M(x - v\Delta t, v, t)$ using definition I and take the minimum $\beta = \min_v \{w(x, v, t + \Delta t)\}$

Off course the different reconstructions are strictly connected to the choice of the macroscopic solver used in point 2.

⁵ S.Deshpande JCP '79, B.Perthame SINUM '90

Macroscopic solvers I

Methods based on discrete velocity model⁶ (HM methods).

Main features

- Representation $f(v) = \tilde{f}(v) + w(v)M(v)$
- Discretize the velocity space.
- Solve the deterministic and stochastic part with a DVM.
- Compact support, equilibrium functions \mathcal{E}_f differ from Maxwellian $\mathcal{E}_f \neq M_f$.
- We need to solve a non linear system for each cell at each time step.
- Time step restrictions from deterministic transport step.

⁶L. Mieussens M³AS '00, L.P. G.Dimarco CMS '07

Macroscopic solvers II

II) Methods based on the full kinetic equation (**BHM methods**).

Main features

- Representation $f(v) = \tilde{f}_R(v) + w_R(v)M(v)$ where $w_R(v) = 0$ for $v \notin [-R, R]^d$.
- Discretize velocity space only in the **central part** $v \in [-R, R]^d$.
- **Tails** are treated by particles.
- Shorter computational time due to time step increase, no need of nonlinear iterations, and to less mesh points in velocity space.
- More fluctuations due to the the presence of the tails.

⁷L.P., G.Dimarco MMS '08, P.Degond, G.Dimarco, L.Mieussens JCP '07

Macroscopic solvers III

III) Methods independent from the fluid solver (FSI methods).

Main features

- Representation $f(v) = \tilde{f}(v) + \beta M(v)$, $\beta = \min_v \{w(v)\}$.
- Solve relaxation in the usual way to get $\beta^r(t)$.
- Solve the transport (equilibrium and nonequilibrium part) with a Monte Carlo method.
- Solve the Euler equations with initial data $U^E(t) = P(\beta^r(t)M(t))$ to get the moments $U^E(t + \Delta t)$. We have

$$P(\beta^r(x - v\Delta t, t)M(x - v\Delta t, v, t)) = U^E(t + \Delta t) + O(\Delta t^2).$$
- Apply a moment matching only to the advected equilibrium particles so that the above equation is satisfied exactly.
- Additional difficulties in the reconstruction since the kinetic information are only available through particles.

⁸L.P., G.Dimarco '08

Accuracy test

Smooth solution in 1D (velocity and space) with periodic boundary conditions. L_1 norm of the errors for temperature respect to different value of the Knudsen number ε (in units of 10^{-2}).

	$\varepsilon = 10^{-2}$	$\varepsilon = 10^{-3}$	$\varepsilon = 10^{-4}$	$\varepsilon = 10^{-5}$	$\varepsilon = 10^{-6}$
MCM	3.2923	4.4354	6.2404	5.7733	6.1142
HM	2.9520	2.7893	2.6305	0.96996	0.2840
HM1	2.8437	2.5110	1.6132	0.6617	0.2053
CHM	1.8196	1.2004	0.5368	0.1310	0.0651
BHM	3.1869	3.0254	2.8536	2.1430	1.8134
BHM1	2.7132	2.6807	2.3756	2.0148	2.1010
BCHM	2.6210	2.3226	2.1498	1.9315	1.8849

$N = 1500$ particles for cell, $v \in [-15, 15]$ for HM schemes, $R = 5$ for *BHM* schemes, $\Delta v = 0.16$ and $\Delta x = 0.05$.

	$\varepsilon = 10^{-2}$	$\varepsilon = 10^{-3}$	$\varepsilon = 5 * 10^{-4}$	$\varepsilon = 10^{-4}$
MCM	6.762	7.611	7.578	7.316
FSI	7.007	6.022	4.500	0.641
FSI1	6.662	4.939	3.773	0.598

$N = 200$ particles for cell $\Delta x = 0.05$.



Computational cost

Smooth solution in 1D (velocity and space) with periodic boundary conditions.

	$\varepsilon = 10^{-2}$	$\varepsilon = 10^{-3}$	$\varepsilon = 10^{-4}$	$\varepsilon = 10^{-5}$
MCM N=1500	23 sec	25 sec	27 sec	26 sec
BHM N=1500	35 sec	25 sec	22 sec	22 sec
BHM1 N=1500	34 sec	20 sec	19 sec	20 sec
BCHM N=1500	15 sec	11 sec	17 sec	21 sec
FSI N=1500	25 sec	22 sec	3 sec	0.6 sec
FSI1 N=1500	18 sec	17 sec	2 sec	0.6 sec
FSI N=500	9 sec	8 sec	0.4 sec	0.3 sec
FSI N=500	7 sec	6 sec	0.4 sec	0.3 sec

Sod test



Comparison of results for ρ for HM and $\varepsilon = 10^{-3}$ ⁹.



⁹ G.Dimarco, L.P. '06



Sod test

Comparison of results for ρ for BHM and $\varepsilon = 10^{-3}$.



BGK equation: flow past an ellipse



Comparison of results for T , DSMC (left), FSI (right).



ε is such that 50% of the solution is represented by particles in HM.

BGK equation: flow past an ellipse


Comparison of results for T , DSMC (left), FSI (right).



The fluid limit $\varepsilon \rightarrow 0$.

Boltzmann equation: 2D channel flow

Comparison of results for ρ (left), T (right), DSMC (left), HM1 (right)¹⁰.



¹⁰ R.Caflich, H.Chen, E.Luo, L.P. AIAA '06

Basic principles

- The basic idea consists in obtaining **reduced variance Monte Carlo** methods forcing particles to match prescribed sets of moments given by the solution of deterministic macroscopic fluid equations⁴.
- These macroscopic models, in order to represent the correct physics for all range of Knudsen numbers include a kinetic correction term, which takes into account departures from thermodynamical equilibrium.
- We will focus on a basic matching technique between the first three moments of the macroscopic and microscopic equations. However, in principle, it is possible to force particles to match also higher order moments, which possibly can further diminish fluctuations.
- The general methodology described in the following is independent from the choice of the collisional kernel (Boltzmann, Fokker-Planck, BGK etc..).

⁴G.Dimarco, P.Degond, L.P., '09

The setting

Consider a kinetic equation of the form

$$\partial_t f + v \cdot \nabla_x f = Q(f, f)$$

The operator $Q(f, f)$ is assumed to satisfy

$$\int_{-\mathbb{R}^3} \phi(v) Q(f, f)(v) dv = 0$$

where $\phi(v) = (1, v, |v|^2)$ are the collision invariants.

We define

$$U = \int_{-\mathbb{R}^3} \phi(v) f(v) dv = (\rho, \rho u, 2E).$$

The HG method

The starting point of the methods is the following **micro-macro decomposition**

$$f(v) = M(v) + g(v).$$

The function $g(v)$ represents the non equilibrium part and it is not strictly positive. Now the moments vector U and $g = f - M$ satisfy the coupled system of equations

$$\partial_t U + \partial_x \int_{\mathbb{R}^3} v f \phi(v) dv + \partial_x \int_{\mathbb{R}^3} v g \phi(v) dv = 0$$

$$\partial_t f + v \partial_x f = Q(f, f).$$

Our scope is to solve the kinetic equation with a Monte Carlo method, and contemporaneously the fluid equation with any type of finite difference or finite volume scheme and than match the resulting moments. Similar decomposition strategies can be used also for low Mach number flows ⁵.

⁵N.Hadjiconstantinou, 05

The HG method

Note that the two systems, with the same initial data, furnish the same results in terms of macroscopic quantities apart from numerical fluctuations.

We summarize the method in the following way



- 1 Solve the kinetic equation and obtain a first set of moments.
- 2 Solve the fluid equation with the preferred finite volume/difference scheme.
- 3 Match the moments of the two models through a transformation of samples values.
- 4 Restart the computation for the next time step.

Step 3 of the above procedure requires great care. If we restrict to moments up to second order then a standard **moment matching** procedure based on a velocity (linear) transformation can be applied.

In the sequel we apply the method to the case of the BGK operator for an unsteady shock problem.



Numerical results: $\varepsilon = 0.1$

Unsteady Shock: Solution at $t = 0.067$ for density. MC method (left), Hydro Guided MC method (right). Knudsen number $\varepsilon = 0.1$.



Numerical results: $\varepsilon = 0.01$

Unsteady Shock: Solution at $t = 0.067$ for density. MC method (left), Hydro Guided MC method (right). Knudsen number $\varepsilon = 0.01$.



Numerical results $\varepsilon = 0.001$

Unsteady Shock: Solution at $t = 0.067$ for density. MC method (left), Hydro Guided MC method (right). Knudsen number $\varepsilon = 0.001$.

