

Approximating Multiples of Strong Rayleigh Random Variables

Thomas M. Liggett, UCLA

Consider a polynomial with positive coefficients

$$f(u) = \sum_{k=0}^n c_k u^k, \quad c_k > 0.$$

It is said to be **Strong Rayleigh** (SR) if all of its roots are real (and hence negative). A random variable X taking values $0, 1, \dots, n$ is SR if its probability generating polynomial (pgf)

$$f(u) = Eu^X = \sum_{k=0}^n P(X = k)u^k$$

is SR.

In this case,

$$f(u) = \prod_{k=0}^n [p_k u + (1 - p_k)]$$

Therefore

$$X =_d \eta_1 + \cdots + \eta_n,$$

where η_i are independent Bernoulli random variables with parameters p_k . If X_n is a sequence of SR random variables, this gives a triangular array

$$X_1 = \eta_{1,1}$$

$$X_2 = \eta_{2,1} + \eta_{2,2}$$

$$X_3 = \eta_{3,1} + \eta_{3,2} + \eta_{3,3}, \dots$$

of Bernoulli random variables with independence in each row. It follows from the Lindeberg-Feller Theorem that if $\text{var}(X_n) \rightarrow \infty$, X_n satisfies the CLT.

Definition A random vector \mathbf{X} is said to be SR if its pgf $f(\mathbf{u}) \neq 0$ whenever $\text{Im}(u_i) > 0$ for all i .

Many natural distributions satisfy SR. But even if one does not know the distribution of \mathbf{X} explicitly, sometimes SR can be verified indirectly.

For example, consider the exclusion process, which is a Markov process on the state space $\{0, 1\}^S$, where S is a countable set. Let $p(x, y)$ be the transition probabilities for a Markov chain on S . Each particle has a rate 1 exponential clock. When the clock at x rings, if there is a particle at x , it tries to move to y with probability $p(x, y)$. If y is occupied, it stays at x ; otherwise it moves to y .

The process is said to be **symmetric** if $p(x, y) = p(y, x)$ for all x, y .

One of many questions about it is the following:

Definition A probability measure on $\{0, 1\}^S$ is said to be negatively associated if $f(\eta)$ and $g(\eta)$ are negatively correlated for all increasing functions f, g that depend on disjoint sets of coordinates.

Problem Is it the case that η_t is negatively associated whenever η_0 is?

Answer No, even in the symmetric case.

However, in

J. Borcea, P. Brändén and T. Liggett. Negative dependence and the geometry of polynomials. *JAMS* **22** (2009) 521–567,

we proved that the symmetric exclusion interacting particle system $\eta_t \in \{0, 1\}^S$ satisfies the following property:

$$\eta_0 \text{ SR} \Rightarrow \eta_t \text{ SR.}$$

Moreover SR implies negative association.

Using this,

T. Liggett. [Distributional limits for the symmetric exclusion process. Stoch. Proc. Appl. **119** \(2009\) 1–15](#)

proved CLT's for the symmetric exclusion process.

In

S. Ghosh, T. Liggett and R. Pemantle. Multivariate CLT follows from strong Rayleigh property. *ANALCO17 (2017)* 139–147,

we raised the question of the extent to which SR implies a multivariate CLT. This is quite different from the univariate case, since the pgf no longer factors, and there is no reason to think that \mathbf{X} can be written as a sum of independent random vectors.

Using a result of Lebowitz, Pittel, Ruelle and Speer, we did prove such a result, but with the assumption $\text{var}(\mathbf{X}_n) \gg n^{\frac{1}{3}}$.

Why do we need a stronger assumption in the multivariate case?

Deducing **multivariate CLT's from univariate CLT's** via the Cramér-Wold device:

$$\mathbf{X}_n \rightarrow_d \mathbf{X} \quad \text{iff} \quad \mathbf{b} \cdot \mathbf{X}_n \rightarrow_d \mathbf{b} \cdot \mathbf{X}$$

for every \mathbf{b} .

This is a simple consequence of the fact that distributional convergence is equivalent to convergence of the characteristic functions (=Fourier transforms).

Problem: If X is SR, bX is not even integer valued, much less SR. Can bX be well approximated by a SR random variable?

Ghosh, Liggett and Pemantle (2017) proved that

if X is SR, then $\lfloor \frac{1}{k}X \rfloor$ is SR.

However, if X is $B(3n, \frac{1}{2})$, then the roots z_i of the pgf of $\lfloor \frac{2}{3}X \rfloor$ satisfy

$$2 \max_i [\operatorname{Im}(z_i)]^2 \geq 9n^2 - 9n - 1.$$

Maybe $\lfloor \frac{j}{k}X \rfloor$, should be written as a sum of independent random variables with more than 2 values....

Theorem. If X is SR, the pgf of $\lfloor \frac{2}{k}X \rfloor$ can be factored into quadratic polynomials with positive coefficients, so $\lfloor \frac{2}{k}X \rfloor$ has the same distribution as the sum of independent random variables taking the values 0,1,2.

Definition f has property P_j if it can be factored into polynomials of degree at most j with positive coefficients.

$$P_1 \iff SR \iff \text{all roots real.}$$

$$P_2 \iff \text{Hurwitz} \iff \text{all roots have negative real part.}$$

P_3 is not a statement about each root. If f is P_3 but not P_2 , each root z with positive real part must be paired with a negative root w so that

$$2\operatorname{Re}(z) < -w < |z|^2/2\operatorname{Re}(z).$$

Theorem (Hermite-Bieler) Write

$$f(u) = \sum_{m=0}^1 u^m h_m(u^2) = h_0(u^2) + u h_1(u^2).$$

Then f is P_2 iff the roots of h_0, h_1 are negative and simple and interlace, with the largest being a root of h_0 .

Definition f has property Q_j if writing

$$f(u) = \sum_{m=0}^{j-1} u^m h_m(u^j),$$

the roots of h_0, h_1, \dots, h_{j-1} are negative and simple and interlace, with the largest being a root of h_0 .

Note that $Q_1 = P_1, Q_2 = P_2$. However, neither implication between Q_3 and P_3 is true.

Location of roots

If f is P_3 , it has no roots in the sector

$$\{z : \operatorname{Re}(z) > 0, (\operatorname{Im}(z))^2 \leq 3(\operatorname{Re}(z))^2\}.$$

If f is Q_3 , it has no roots on

$$\{z : \operatorname{Re}(z) > 0, (\operatorname{Im}(z))^2 = 3(\operatorname{Re}(z))^2\}.$$

Theorem If X is SR, then $\lfloor \frac{j}{k} X \rfloor$ is Q_j .

Corollary If X is SR, then $\lfloor \frac{2}{k} X \rfloor$ is P_2 .

Conjecture If X is SR, then $\lfloor \frac{3}{4} X \rfloor$ is P_3 .

This is true if $X \leq 6$. If X is $B(40, p)$ with $p = \frac{1}{8}, \frac{1}{4}$ or $\frac{1}{2}$, the pgf of $\lfloor \frac{3}{4} X \rfloor$ has 10 real roots

$$w_{10} < w_9 < \cdots < w_1 < 0$$

and 10 conjugate pairs of roots

$$z_1, \bar{z}_1, \dots, z_{10}, \bar{z}_{10}$$

with $0 < \operatorname{Re}(z_1) < \cdots < \operatorname{Re}(z_{10})$. With this ordering, $(u - w_i)(u - z_i)(u - \bar{z}_i)$ has positive coefficients for each $1 \leq i \leq 10$.

If X is $B(21, \frac{1}{2})$, $\lfloor \frac{3}{5}X \rfloor$ is not P_3 . However, it is almost P_3 in the sense that its pgf is the product 4 cubics, only one of which has a negative coefficient:

$$22(.00031 + .021u - .0058u^2 + u^3)(.12 + .43u + .14u^2 + u^3) \\ \cdot (8.96 + 5.88u + .92u^2 + u^3)(2993 + 317u + 8.49u^2 + u^3).$$

The same pattern occurs if X is $B(n, \frac{1}{2})$ for $n = 25, 35, 50$.

Proposition If U, V are nonnegative integer valued random variables whose pgf's satisfy

$$Eu^U = Eu^V (au^3 + bu^2 + cu + d)$$

with $d \geq 0, c + d \geq 0, b + c + d \geq 0$, then there is a coupling so that $U \leq V + 3$ and $E(V + 3 - U) = b + 2c + 3d$.

The underlying fact that our results depend on involves polynomials with interlacing roots:

Theorem (Ghosh, Liggett, Pemantle). Let f be the pgf of a SR X taking values $0, 1, \dots, n$, which is a polynomial of degree n with all negative roots. Write

$$f(x) = \sum_{i=0}^{k-1} x^i g_i(x^k),$$

where g_i is a polynomial of degree $\lfloor \frac{n-i}{k} \rfloor$. Then g_0, g_1, \dots, g_{k-1} have interlacing, negative simple roots, with the largest being a root of g_0 .

The proof is by induction on the degree of f .

For the induction argument, write $F(x) = (x + r)f(x)$ with $r > 0$, where f has degree n and F has degree $n + 1$. Consider the corresponding decomposition for F :

$$F(x) = \sum_{i=0}^{k-1} x^i G_i(x^k).$$

Then

$$G_i(y) = rg_i(y) + \begin{cases} yg_{k-1}(y) & \text{if } i = 0; \\ g_{i-1}(y) & \text{if } i \geq 1. \end{cases}$$

Let the roots of the g_i 's be $\cdots < s_4 < s_3 < s_2 < s_1 < s_0 < 0$.

Then for $k = 3$, for example, the following explains the proof.

$$\begin{pmatrix} & \cdots & s_6 & s_5 & s_4 & s_3 & s_2 & s_1 & s_0 & 0 \\ g_0 & \cdots & 0 & + & + & 0 & - & - & 0 & + \\ g_1 & \cdots & + & + & 0 & - & - & 0 & + & + \\ g_2 & \cdots & + & 0 & - & - & 0 & + & + & + \\ G_0 & \cdots & - & + & + & + & - & - & - & + \\ G_1 & \cdots & + & + & + & - & - & - & + & + \\ G_2 & \cdots & + & + & - & - & - & + & + & + \end{pmatrix} .$$

So, G_0 has a root in $\dots, (s_3, s_2), (s_0, 0)$, G_1 has a root in $\dots, (s_4, s_3), (s_1, s_0)$, and G_2 has a root in $\dots, (s_5, s_4), (s_2, s_1)$.

The proof that $\lfloor \frac{j}{k} X \rfloor$ is Q_j if X is SR is similar. The h_i 's in the definition of property Q_j are

$$h_i(u) = \sum_{ik \leq mj < (i+1)k} g_m(u).$$

For $j = 4, k = 7$

$$h_0 = g_0 + g_1, \quad h_1 = g_2 + g_3, \quad h_2 = g_4 + g_5, \quad h_3 = g_6.$$

The proof is described in the following form:

$$\begin{pmatrix}
 & \cdots & s_7 & s_6 & s_5 & s_4 & s_3 & s_2 & s_1 & s_0 \\
 g_0 & \cdots & 0 & - & - & - & - & - & - & 0 \\
 g_1 & \cdots & - & - & - & - & - & - & 0 & + \\
 g_2 & \cdots & - & - & - & - & - & 0 & + & + \\
 g_3 & \cdots & - & - & - & - & 0 & + & + & + \\
 g_4 & \cdots & - & - & - & 0 & + & + & + & + \\
 g_5 & \cdots & - & - & 0 & + & + & + & + & + \\
 g_6 & \cdots & - & 0 & + & + & + & + & + & + \\
 h_0 & \cdots & - & - & - & - & - & - & - & + \\
 h_1 & \cdots & - & - & - & - & - & + & + & + \\
 h_2 & \cdots & - & - & - & + & + & + & + & + \\
 h_3 & \cdots & - & 0 & + & + & + & + & + & +
 \end{pmatrix}$$