

Gravitational allocation to Poisson points

Yuval Peres

joint work with

Sourav Chatterjee

Ron Peled

Dan Romik

Allocation rules

- ▶ Let Ξ be a discrete subset of \mathbb{R}^d .

Allocation rules

- ▶ Let Ξ be a discrete subset of \mathbb{R}^d .
- ▶ An **allocation** (of Lebesgue measure to Ξ) is a measurable function $\psi : \mathbb{R}^d \rightarrow \Xi \cup \{\infty\}$ that satisfies

$$\begin{aligned}\text{Vol}(\psi^{-1}(\infty)) &= 0, \\ \text{Vol}(\psi^{-1}(z)) &= 1, \quad z \in \Xi,\end{aligned}$$

where $\text{Vol}(\cdot)$ is Lebesgue measure in \mathbb{R}^d .

Allocation rules


- ▶ Let Ξ be a discrete subset of \mathbb{R}^d .
- ▶ An **allocation** (of Lebesgue measure to Ξ) is a measurable function $\psi : \mathbb{R}^d \rightarrow \Xi \cup \{\infty\}$ that satisfies

$$\begin{aligned}\text{Vol}(\psi^{-1}(\infty)) &= 0, \\ \text{Vol}(\psi^{-1}(z)) &= 1, \quad z \in \Xi,\end{aligned}$$


where $\text{Vol}(\cdot)$ is Lebesgue measure in \mathbb{R}^d .

- ▶ For $x \in \Xi$, we call $\psi^{-1}(z)$ the **cell allocated to z** .

Examples



(a)



(b)

Figure: (a) The two-dimensional stable marriage allocation for a Poisson process (picture due to Alexander E. Holroyd). (b) The gradient flow allocation (picture due to Manjunath Krishnapur).

Allocation rules contd.

- ▶ Let Z be a translation-invariant simple point process in \mathbb{R}^d with unit intensity.

Allocation rules contd.

- ▶ Let Z be a translation-invariant simple point process in \mathbb{R}^d with unit intensity.
- ▶ An **invariant allocation rule** (of Lebesgue measure to Z) is a measurable mapping $Z \rightarrow \psi_Z$ such that:

Allocation rules contd.

- ▶ Let Z be a translation-invariant simple point process in \mathbb{R}^d with unit intensity.
- ▶ An **invariant allocation rule** (of Lebesgue measure to Z) is a measurable mapping $Z \rightarrow \psi_Z$ such that:
 1. a.s. ψ_Z is an allocation of Lebesgue measure to Z , and

Allocation rules contd.

- ▶ Let Z be a translation-invariant simple point process in \mathbb{R}^d with unit intensity.
- ▶ An **invariant allocation rule** (of Lebesgue measure to Z) is a measurable mapping $Z \rightarrow \psi_Z$ such that:
 1. a.s. ψ_Z is an allocation of Lebesgue measure to Z , and
 2. the mapping $Z \rightarrow \psi_Z$ is translation-equivariant, i.e.

$$\psi_{Z+x}(y) \equiv \psi_Z(y) + x.$$

Allocation rules contd.

- ▶ Let Z be a translation-invariant simple point process in \mathbb{R}^d with unit intensity.
- ▶ An **invariant allocation rule** (of Lebesgue measure to Z) is a measurable mapping $Z \rightarrow \psi_Z$ such that:
 1. a.s. ψ_Z is an allocation of Lebesgue measure to Z , and
 2. the mapping $Z \rightarrow \psi_Z$ is translation-equivariant, i.e.

$$\psi_{Z+x}(y) \equiv \psi_Z(y) + x.$$

- ▶ If a.s. all the cells are bounded, one can consider the **allocation diameter**

$$X = \text{diam}(\psi_Z^{-1}(\psi_Z(0))).$$

Allocation rules contd.

- ▶ Let Z be a translation-invariant simple point process in \mathbb{R}^d with unit intensity.
- ▶ An **invariant allocation rule** (of Lebesgue measure to Z) is a measurable mapping $Z \rightarrow \psi_Z$ such that:
 1. a.s. ψ_Z is an allocation of Lebesgue measure to Z , and
 2. the mapping $Z \rightarrow \psi_Z$ is translation-equivariant, i.e.

$$\psi_{Z+x}(y) \equiv \psi_Z(y) + x.$$

- ▶ If a.s. all the cells are bounded, one can consider the **allocation diameter**

$$X = \text{diam}(\psi_Z^{-1}(\psi_Z(0))).$$

- ▶ One object of interest: The rate of decay of the tail $\mathbb{P}(X > R)$.

- ▶ Let Z be a translation-invariant simple point process on \mathbb{R}^d with unit intensity.

Extra head rules

- ▶ Let Z be a translation-invariant simple point process on \mathbb{R}^d with unit intensity.
- ▶ An extra head rule for Z is a random variable T coupled with Z so that

Extra head rules

- ▶ Let Z be a translation-invariant simple point process on \mathbb{R}^d with unit intensity.
- ▶ An extra head rule for Z is a random variable T coupled with Z so that
 1. a.s. $T \in Z$, and

Extra head rules

- ▶ Let Z be a translation-invariant simple point process on \mathbb{R}^d with unit intensity.
- ▶ An extra head rule for Z is a random variable T coupled with Z so that
 1. a.s. $T \in Z$, and
 2. the random set $Z - T$ has the same distribution as Z conditioned to have a point at 0.

Extra head rules

- ▶ Let Z be a translation-invariant simple point process on \mathbb{R}^d with unit intensity.
- ▶ An extra head rule for Z is a random variable T coupled with Z so that
 1. a.s. $T \in Z$, and
 2. the random set $Z - T$ has the same distribution as Z conditioned to have a point at 0.
- ▶ The extra head rule is said to be non-randomized if T is measurable with respect to Z .

Extra head rules

- ▶ Let Z be a translation-invariant simple point process on \mathbb{R}^d with unit intensity.
- ▶ An extra head rule for Z is a random variable T coupled with Z so that
 1. a.s. $T \in Z$, and
 2. the random set $Z - T$ has the same distribution as Z conditioned to have a point at 0.
- ▶ The extra head rule is said to be non-randomized if T is measurable with respect to Z .
- ▶ Holroyd & Peres showed that if ψ_Z is an invariant allocation rule then $T = T_Z = \psi_Z(0)$ is a non-randomized extra head rule,

Extra head rules

- ▶ Let Z be a translation-invariant simple point process on \mathbb{R}^d with unit intensity.
- ▶ An extra head rule for Z is a random variable T coupled with Z so that
 1. a.s. $T \in Z$, and
 2. the random set $Z - T$ has the same distribution as Z conditioned to have a point at 0.
- ▶ The extra head rule is said to be non-randomized if T is measurable with respect to Z .
- ▶ Holroyd & Peres showed that if ψ_Z is an invariant allocation rule then $T = T_Z = \psi_Z(0)$ is a non-randomized extra head rule, and conversely, given a non-randomized extra head rule T_Z , the mapping $\psi_Z(x) = x + T_{Z-x}$ is an invariant allocation rule.

Allocation to Poisson points: Existing results

- ▶ Talagrand: Randomized allocations for $d \geq 3$. Holroyd & Peres used this to construct an allocation with $\mathbb{P}(\psi_Z(0) > R) \leq C \exp(-cR^d)$, where $\psi_Z(0)$ is the typical allocation distance.

Allocation to Poisson points: Existing results

- ▶ Talagrand: Randomized allocations for $d \geq 3$. Holroyd & Peres used this to construct an allocation with $\mathbb{P}(\psi_Z(0) > R) \leq C \exp(-cR^d)$, where $\psi_Z(0)$ is the typical allocation distance.
- ▶ Holroyd & Peres: If $d = 1, 2$ and Z is a standard Poisson point process of unit intensity in \mathbb{R}^d , then the allocation diameter of any invariant rule X satisfies $\mathbb{E}X^{d/2} = \infty$.

Allocation to Poisson points: Existing results

- ▶ Talagrand: Randomized allocations for $d \geq 3$. Holroyd & Peres used this to construct an allocation with $\mathbb{P}(\psi_Z(0) > R) \leq C \exp(-cR^d)$, where $\psi_Z(0)$ is the typical allocation distance.
- ▶ Holroyd & Peres: If $d = 1, 2$ and Z is a standard Poisson point process of unit intensity in \mathbb{R}^d , then the allocation diameter of any invariant rule X satisfies $\mathbb{E}X^{d/2} = \infty$.
- ▶ Hoffman, Holroyd & Peres: For arbitrary translation-invariant point process in \mathbb{R}^d , $d \geq 1$, constructed the **stable marriage allocation**.

Allocation to Poisson points: Existing results

- ▶ Talagrand: Randomized allocations for $d \geq 3$. Holroyd & Peres used this to construct an allocation with $\mathbb{P}(\psi_Z(0) > R) \leq C \exp(-cR^d)$, where $\psi_Z(0)$ is the typical allocation distance.
- ▶ Holroyd & Peres: If $d = 1, 2$ and Z is a standard Poisson point process of unit intensity in \mathbb{R}^d , then the allocation diameter of any invariant rule X satisfies $\mathbb{E}X^{d/2} = \infty$.
- ▶ Hoffman, Holroyd & Peres: For arbitrary translation-invariant point process in \mathbb{R}^d , $d \geq 1$, constructed the **stable marriage allocation**.
- ▶ In the stable marriage allocation, a.s.
 1. all the cells are bounded and contain their owners,

Allocation to Poisson points: Existing results

- ▶ Talagrand: Randomized allocations for $d \geq 3$. Holroyd & Peres used this to construct an allocation with $\mathbb{P}(\psi_Z(0) > R) \leq C \exp(-cR^d)$, where $\psi_Z(0)$ is the typical allocation distance.
- ▶ Holroyd & Peres: If $d = 1, 2$ and Z is a standard Poisson point process of unit intensity in \mathbb{R}^d , then the allocation diameter of any invariant rule X satisfies $\mathbb{E}X^{d/2} = \infty$.
- ▶ Hoffman, Holroyd & Peres: For arbitrary translation-invariant point process in \mathbb{R}^d , $d \geq 1$, constructed the **stable marriage allocation**.
- ▶ In the stable marriage allocation, a.s.
 1. all the cells are bounded and contain their owners,
 2. but not all are connected,

Allocation to Poisson points: Existing results

- ▶ Talagrand: Randomized allocations for $d \geq 3$. Holroyd & Peres used this to construct an allocation with $\mathbb{P}(\psi_Z(0) > R) \leq C \exp(-cR^d)$, where $\psi_Z(0)$ is the typical allocation distance.
- ▶ Holroyd & Peres: If $d = 1, 2$ and Z is a standard Poisson point process of unit intensity in \mathbb{R}^d , then the allocation diameter of any invariant rule X satisfies $\mathbb{E}X^{d/2} = \infty$.
- ▶ Hoffman, Holroyd & Peres: For arbitrary translation-invariant point process in \mathbb{R}^d , $d \geq 1$, constructed the **stable marriage allocation**.
- ▶ In the stable marriage allocation, a.s.
 1. all the cells are bounded and contain their owners,
 2. but not all are connected,
 3. and when Z is a Poisson point process the allocation diameter X satisfies $\mathbb{E}X^d = \infty$.

Picture of a stable marriage allocation




Figure: The 2-diml. stable marriage allocation for a Poisson process

Construction: Each star (point of the process) grows a ball at unit rate and captures all the sites it reaches first, until it is sated (has obtained volume 1).

- Allocation to the zeros of the Gaussian Entire Function (GEF)

$$f(z) = \sum_{n=0}^{\infty} \xi_n \frac{z^n}{\sqrt{n!}}, \quad z \in \mathbb{C},$$

where $(\xi_n)_{n=1}^{\infty}$ are i.i.d. standard complex gaussian random variables.

- Allocation to the zeros of the Gaussian Entire Function (GEF)

$$f(z) = \sum_{n=0}^{\infty} \xi_n \frac{z^n}{\sqrt{n!}}, \quad z \in \mathbb{C},$$

where $(\xi_n)_{n=1}^{\infty}$ are i.i.d. standard complex gaussian random variables.

- Cell of each zero z is defined as the **basin of attraction** of z with respect to the flow induced by the random planar vector field $F(z) = z - (\nabla \log |f|)(z)$. (See next slide for a picture of the corresponding potential $U(z) = \log |f|(z) - \frac{1}{2}|z|^2$.)

- Allocation to the zeros of the Gaussian Entire Function (GEF)

$$f(z) = \sum_{n=0}^{\infty} \xi_n \frac{z^n}{\sqrt{n!}}, \quad z \in \mathbb{C},$$

where $(\xi_n)_{n=1}^{\infty}$ are i.i.d. standard complex gaussian random variables.

- Cell of each zero z is defined as the **basin of attraction** of z with respect to the flow induced by the random planar vector field $F(z) = z - (\nabla \log |f|)(z)$. (See next slide for a picture of the corresponding potential $U(z) = \log |f|(z) - \frac{1}{2}|z|^2$.)
- Construction due to Nazarov, Sodin & Volberg based on an idea suggested by Tsirelson.

Picture of planar potential




Figure: The random planar potential (courtesy of Manjunath Krishnapur).

Picture of a gradient flow allocation




Figure: The gradient flow allocation (courtesy of Manjunath Krishnapur).

Nazarov, Sodin & Volberg showed that the cells are connected, a.s. bounded, and there exist absolute constants $C, c > 0$ such that the allocation diameter X satisfies

$$ce^{-CR(\log R)^{3/2}} \leq \mathbb{P}(X \geq R) \leq Ce^{-cR(\log R)^{3/2}}, \quad R > 1.$$

Why equal area in each basin?

- ▶ Take a basin of attraction $B(z)$, and a point $x \in \partial B(z)$.

Why equal area in each basin?

- ▶ Take a basin of attraction $B(z)$, and a point $x \in \partial B(z)$.
- ▶ If \mathbf{n} is the outward-pointing normal vector at x , then by the definition of the basin of attraction, $F(x) \cdot \mathbf{n} = 0$.

Why equal area in each basin?

- ▶ Take a basin of attraction $B(z)$, and a point $x \in \partial B(z)$.
- ▶ If \mathbf{n} is the outward-pointing normal vector at x , then by the definition of the basin of attraction, $F(x) \cdot \mathbf{n} = 0$.
- ▶ Thus, the oriented surface integral

$$\int_{\partial B(z)} F(x) \cdot \mathbf{n} \, dS = 0.$$

Why equal area in each basin?

- ▶ Take a basin of attraction $B(z)$, and a point $x \in \partial B(z)$.
- ▶ If \mathbf{n} is the outward-pointing normal vector at x , then by the definition of the basin of attraction, $F(x) \cdot \mathbf{n} = 0$.
- ▶ Thus, the oriented surface integral

$$\int_{\partial B(z)} F(x) \cdot \mathbf{n} \, dS = 0.$$

- ▶ Now

$$\operatorname{div}(F) = 2 - 2\pi \sum_{i=1}^{\infty} \delta_{z_i},$$

where $(z_i)_i$ is the set of zeros.

Why equal area in each basin?

- ▶ Take a basin of attraction $B(z)$, and a point $x \in \partial B(z)$.
- ▶ If \mathbf{n} is the outward-pointing normal vector at x , then by the definition of the basin of attraction, $F(x) \cdot \mathbf{n} = 0$.
- ▶ Thus, the oriented surface integral

$$\int_{\partial B(z)} F(x) \cdot \mathbf{n} \, dS = 0.$$

- ▶ Now

$$\operatorname{div}(F) = 2 - 2\pi \sum_{i=1}^{\infty} \delta_{z_i},$$

where $(z_i)_i$ is the set of zeros.

- ▶ Thus, by the divergence theorem,

$$\int_{\partial B(z)} F(x) \cdot \mathbf{n} \, dS = \int_{B(z)} \operatorname{div}(F) \, dx = 2\operatorname{Vol}(B(z)) - 2\pi.$$

Why equal area in each basin?

- ▶ Take a basin of attraction $B(z)$, and a point $x \in \partial B(z)$.
- ▶ If \mathbf{n} is the outward-pointing normal vector at x , then by the definition of the basin of attraction, $F(x) \cdot \mathbf{n} = 0$.
- ▶ Thus, the oriented surface integral

$$\int_{\partial B(z)} F(x) \cdot \mathbf{n} \, dS = 0.$$

- ▶ Now

$$\operatorname{div}(F) = 2 - 2\pi \sum_{i=1}^{\infty} \delta_{z_i},$$

where $(z_i)_i$ is the set of zeros.

- ▶ Thus, by the divergence theorem,

$$\int_{\partial B(z)} F(x) \cdot \mathbf{n} \, dS = \int_{B(z)} \operatorname{div}(F) \, dx = 2\operatorname{Vol}(B(z)) - 2\pi.$$

- ▶ Combining, we get $\operatorname{Vol}(B(z)) = \pi$.

Invariance of the travel time

- Fix $\epsilon > 0$, a zero z , and for each $t \geq 0$, let

$$D(t) = \{x = \Gamma(0) : |\Gamma(t) - z| < \epsilon\}$$

be the ‘region covered in time t ’.

Invariance of the travel time

- Fix $\epsilon > 0$, a zero z , and for each $t \geq 0$, let

$$D(t) = \{x = \Gamma(0) : |\Gamma(t) - z| < \epsilon\}$$

be the ‘region covered in time t ’.

- Let $A(t)$ be the area of $B(z) \setminus D(t)$. By Liouville’s theorem and the fact that $\operatorname{div}(F) = -2$ outside $(z_i)_i$, it follows that

$$\frac{dA}{dt} = -2A(t).$$

Invariance of the travel time

- Fix $\epsilon > 0$, a zero z , and for each $t \geq 0$, let

$$D(t) = \{x = \Gamma(0) : |\Gamma(t) - z| < \epsilon\}$$

be the 'region covered in time t '.

- Let $A(t)$ be the area of $B(z) \setminus D(t)$. By Liouville's theorem and the fact that $\operatorname{div}(F) = -2$ outside $(z_i)_i$, it follows that

$$\frac{dA}{dt} = -2A(t).$$

- Gives alternative proof of the equal area theorem. Also, shows that if τ_x is the 'time' taken by $x \in B(z)$ to 'roll down' to z , then $\mathbb{P}(\tau_x > t) = e^{-2t}$.

Invariance of the travel time

- Fix $\epsilon > 0$, a zero z , and for each $t \geq 0$, let

$$D(t) = \{x = \Gamma(0) : |\Gamma(t) - z| < \epsilon\}$$

be the ‘region covered in time t ’.

- Let $A(t)$ be the area of $B(z) \setminus D(t)$. By Liouville’s theorem and the fact that $\operatorname{div}(F) = -2$ outside $(z_i)_i$, it follows that

$$\frac{dA}{dt} = -2A(t).$$

- Gives alternative proof of the equal area theorem. Also, shows that if τ_x is the ‘time’ taken by $x \in B(z)$ to ‘roll down’ to z , then $\mathbb{P}(\tau_x > t) = e^{-2t}$.
- In fact, a stronger version holds, where we condition on the allocation and choose x uniformly in one cell.

Gravitational force field due to Poisson points

- ▶ Let \mathcal{Z} be a standard Poisson process in \mathbb{R}^d .

Gravitational force field due to Poisson points

- ▶ Let \mathcal{Z} be a standard Poisson process in \mathbb{R}^d .
- ▶ Consider the random vector field $F : \mathbb{R}^d \rightarrow \mathbb{R}^d$ defined by

$$F(x) = \sum_{z \in \mathcal{Z}, |z-x| \uparrow} \frac{z-x}{|z-x|^d}, \quad (1)$$

where the summands are arranged in order of increasing distance from x .

Gravitational force field due to Poisson points

- ▶ Let \mathcal{Z} be a standard Poisson process in \mathbb{R}^d .
- ▶ Consider the random vector field $F : \mathbb{R}^d \rightarrow \mathbb{R}^d$ defined by

$$F(x) = \sum_{z \in \mathcal{Z}, |z-x| \uparrow} \frac{z-x}{|z-x|^d}, \quad (1)$$

where the summands are arranged in order of increasing distance from x .

- ▶ First investigated in work of S. Chandrasekhar. Later work by Heath & Shepp.

Convergence of the force at the origin

- ▶ Since $\mathcal{Z} = (z_i)_i$ is a Poisson process in \mathbb{R}^d with unit intensity, it follows that $(|z_i|^d)_i$ is a Poisson process in \mathbb{R} with intensity κ_d , where κ_d is the volume of the unit ball in \mathbb{R}^d .

Convergence of the force at the origin

- ▶ Since $\mathcal{Z} = (z_i)_i$ is a Poisson process in \mathbb{R}^d with unit intensity, it follows that $(|z_i|^d)_i$ is a Poisson process in \mathbb{R} with intensity κ_d , where κ_d is the volume of the unit ball in \mathbb{R}^d .
- ▶ Thus, we have

$$\frac{|z_i|}{i^{1/d}} \xrightarrow{i \rightarrow \infty} \kappa_d^{-1/d} \text{ a.s.}$$

Convergence of the force at the origin

- ▶ Since $\mathcal{Z} = (z_i)_i$ is a Poisson process in \mathbb{R}^d with unit intensity, it follows that $(|z_i|^d)_i$ is a Poisson process in \mathbb{R} with intensity κ_d , where κ_d is the volume of the unit ball in \mathbb{R}^d .
- ▶ Thus, we have

$$\frac{|z_i|}{i^{1/d}} \xrightarrow{i \rightarrow \infty} \kappa_d^{-1/d} \text{ a.s.}$$

- ▶ If we condition on $(|z_i|)_i$, then each z_i is distributed uniformly on the sphere of radius $|z_i|$ around the origin, and they are independent.

Convergence of the force at the origin

- ▶ Since $\mathcal{Z} = (z_i)_i$ is a Poisson process in \mathbb{R}^d with unit intensity, it follows that $(|z_i|^d)_i$ is a Poisson process in \mathbb{R} with intensity κ_d , where κ_d is the volume of the unit ball in \mathbb{R}^d .
- ▶ Thus, we have

$$\frac{|z_i|}{i^{1/d}} \xrightarrow{i \rightarrow \infty} \kappa_d^{-1/d} \text{ a.s.}$$

- ▶ If we condition on $(|z_i|)_i$, then each z_i is distributed uniformly on the sphere of radius $|z_i|$ around the origin, and they are independent.
- ▶ For each i , $z_i/|z_i|^d$ has conditional mean 0 and variance bounded by $O(|z_i|^{-2(d-1)}) = O(i^{-2(d-1)/d})$.

Convergence of the force at the origin

- ▶ Since $\mathcal{Z} = (z_i)_i$ is a Poisson process in \mathbb{R}^d with unit intensity, it follows that $(|z_i|^d)_i$ is a Poisson process in \mathbb{R} with intensity κ_d , where κ_d is the volume of the unit ball in \mathbb{R}^d .
- ▶ Thus, we have

$$\frac{|z_i|}{i^{1/d}} \xrightarrow{i \rightarrow \infty} \kappa_d^{-1/d} \text{ a.s.}$$

- ▶ If we condition on $(|z_i|)_i$, then each z_i is distributed uniformly on the sphere of radius $|z_i|$ around the origin, and they are independent.
- ▶ For each i , $z_i/|z_i|^d$ has conditional mean 0 and variance bounded by $O(|z_i|^{-2(d-1)}) = O(i^{-2(d-1)/d})$.
- ▶ Thus, a.s. convergence of $F(0)$ follows from the Kolmogorov 3-series theorem. By stationarity, same is true for every $F(x)$.

Convergence of the force at the origin

- ▶ Since $\mathcal{Z} = (z_i)_i$ is a Poisson process in \mathbb{R}^d with unit intensity, it follows that $(|z_i|^d)_i$ is a Poisson process in \mathbb{R} with intensity κ_d , where κ_d is the volume of the unit ball in \mathbb{R}^d .
- ▶ Thus, we have

$$\frac{|z_i|}{i^{1/d}} \xrightarrow{i \rightarrow \infty} \kappa_d^{-1/d} \text{ a.s.}$$

- ▶ If we condition on $(|z_i|)_i$, then each z_i is distributed uniformly on the sphere of radius $|z_i|$ around the origin, and they are independent.
- ▶ For each i , $z_i/|z_i|^d$ has conditional mean 0 and variance bounded by $O(|z_i|^{-2(d-1)}) = O(i^{-2(d-1)/d})$.
- ▶ Thus, a.s. convergence of $F(0)$ follows from the Kolmogorov 3-series theorem. By stationarity, same is true for every $F(x)$.
- ▶ In our paper we have shown that a.s. the series converges everywhere on $\mathbb{R}^d \setminus \mathcal{Z}$ to give a translation-invariant continuously differentiable random function.

Distribution of $F(0)$

- ▶ Each component of the vector $F(0)$ has a stable distribution whose parameters can be explicitly computed (special case by Chandrasekhar, general proof by Heath & Shepp). We give a sketch below.

Distribution of $F(0)$

- ▶ Each component of the vector $F(0)$ has a stable distribution whose parameters can be explicitly computed (special case by Chandrasekhar, general proof by Heath & Shepp). We give a sketch below.
- ▶ If F_1, \dots, F_n are i.i.d. copies of the force F , then their sum is the force exerted by the union of n copies of the poisson process, which is Poisson scaled by $n^{-1/d}$.

Distribution of $F(0)$

- ▶ Each component of the vector $F(0)$ has a stable distribution whose parameters can be explicitly computed (special case by Chandrasekhar, general proof by Heath & Shepp). We give a sketch below.
- ▶ If F_1, \dots, F_n are i.i.d. copies of the force F , then their sum is the force exerted by the union of n copies of the poisson process, which is Poisson scaled by $n^{-1/d}$.
- ▶ Thus, $F_1(0) + \dots + F_n(0)$ has same law as $n^{(d-1)/d} F(0)$.

Distribution of $F(0)$

- ▶ Each component of the vector $F(0)$ has a stable distribution whose parameters can be explicitly computed (special case by Chandrasekhar, general proof by Heath & Shepp). We give a sketch below.
- ▶ If F_1, \dots, F_n are i.i.d. copies of the force F , then their sum is the force exerted by the union of n copies of the poisson process, which is Poisson scaled by $n^{-1/d}$.
- ▶ Thus, $F_1(0) + \dots + F_n(0)$ has same law as $n^{(d-1)/d} F(0)$.
- ▶ It follows that $F(0)$ is $d/(d-1)$ stable.

A rearrangement identity

Lemma

For each $u, x \in \mathbb{R}^d$, let

$$G^{\{u\}}(x) = \sum_{|z_i - u| \uparrow} \frac{z_i - x}{|z_i - x|^d}.$$

Then for any $x, u, v \in \mathbb{R}^d$ we have $G^{\{u\}}(x) - G^{\{v\}}(x) = \kappa_d(u - v)$ a.s., where κ_d is the volume of the unit ball in \mathbb{R}^d .

A rearrangement identity

Lemma

For each $u, x \in \mathbb{R}^d$, let

$$G^{\{u\}}(x) = \sum_{|z_i - u| \uparrow} \frac{z_i - x}{|z_i - x|^d}.$$

Then for any $x, u, v \in \mathbb{R}^d$ we have $G^{\{u\}}(x) - G^{\{v\}}(x) = \kappa_d(u - v)$ a.s., where κ_d is the volume of the unit ball in \mathbb{R}^d .

Proof:

- ▶ If $N_{u,x}$ is the number of stars in the ball $B(u, |u - x|)$, then

$$\mathbb{E}\left[G^{\{u\}}(x) \middle| N_{u,x}\right] = N_{u,x} \cdot \frac{u - x}{|u - x|^d}.$$

(Follows from a well-known physics principle.) Thus,
 $\mathbb{E}(G^{\{u\}}(x) - G^{\{v\}}(x)) = \kappa_d(u - v).$

Contd. on next slide...

Rearrangement identity contd.


- Now, let $R > 0$ be large, and consider the truncated series

$$G_R^{\{u\}}(x) = \sum_{|z_i - u| < R} \frac{z_i - x}{|z_i - x|^d}.$$

We show that $\text{Var}(G_R^{\{u\}}(x) - G_R^{\{v\}}(x)) \rightarrow 0$ as $R \rightarrow \infty$. This suffices to complete the proof of the rearrangement identity.

Contd. on next slide...

Rearrangement identity contd.



- ▶ Since $G_R^{\{u\}}(x) - G_R^{\{v\}}(x)$ is the sum of independent contributions from the E_j 's, the variance can be easily bounded.

Gravitational allocation

- ▶ Consider now the integral curves $\Gamma(t)$ of the vector field F , that is, solutions of the equation

$$\dot{\Gamma}(t) = F(\Gamma(t)).$$

We call these curves the **gravitational flow curves**.

Gravitational allocation

- ▶ Consider now the integral curves $\Gamma(t)$ of the vector field F , that is, solutions of the equation

$$\dot{\Gamma}(t) = F(\Gamma(t)).$$

We call these curves the **gravitational flow curves**.

- ▶ Denote by Γ_x the integral curve with initial condition $\Gamma_x(0) = x$.

Gravitational allocation

- ▶ Consider now the integral curves $\Gamma(t)$ of the vector field F , that is, solutions of the equation

$$\dot{\Gamma}(t) = F(\Gamma(t)).$$

We call these curves the **gravitational flow curves**.

- ▶ Denote by Γ_x the integral curve with initial condition $\Gamma_x(0) = x$.
- ▶ To each center $z \in \mathcal{Z}$, define its **basin of attraction**

$$B(z) = \{x \in \mathbb{R}^d \setminus \mathcal{Z} \mid \Gamma_x(t) \text{ ends at } z\} \cup \{z\}.$$

Gravitational allocation

- ▶ Consider now the integral curves $\Gamma(t)$ of the vector field F , that is, solutions of the equation

$$\dot{\Gamma}(t) = F(\Gamma(t)).$$

We call these curves the **gravitational flow curves**.

- ▶ Denote by Γ_x the integral curve with initial condition $\Gamma_x(0) = x$.
- ▶ To each center $z \in \mathcal{Z}$, define its **basin of attraction**

$$B(z) = \{x \in \mathbb{R}^d \setminus \mathcal{Z} \mid \Gamma_x(t) \text{ ends at } z\} \cup \{z\}.$$

- ▶ Define the **gravitational allocation rule**

$$\psi_{\mathcal{Z}}(x) = \begin{cases} z & x \in B(z) \text{ for } z \in \mathcal{Z}, \\ \infty & x \notin \bigcup_{z \in \mathcal{Z}} B(z). \end{cases}$$

Theorem

The mapping $\mathcal{Z} \rightarrow \psi_{\mathcal{Z}}$ is an allocation rule of Lebesgue measure to the Poisson point process \mathcal{Z} . Almost surely all the cells $\psi^{-1}(z)$ are bounded. The allocation diameter $X = \text{diam}(\psi^{-1}(\psi(0)))$ satisfies the following tail bounds: In dimensions 4 and higher, we have

$$\mathbb{P}(X > R) \leq C_1 \exp \left[-c_2 R (\log R)^{\frac{d-2}{d}} \right] \quad (2)$$

for some constants $C_1, c_2 > 0$ (depending on the dimension d) and all positive R . In dimension 3, for any $\alpha > 0$ there exist constants $C_1, c_2 > 0$ (depending on α) such that for all $R > 0$ we have

$$\mathbb{P}(X > R) \leq C_1 \exp \left[-c_2 \frac{R}{(\log R)^{\frac{4}{3} + \alpha}} \right]. \quad (3)$$

Picture of gravitational allocation




Figure: Simulation of a cell in 3-dimensional gravitational allocation

Sketch of the proof

- ▶ We closely follow a technique introduced in an earlier version of the paper of Nazarov, Sodin & Volberg, but several new ideas are required to carry out the steps.

Sketch of the proof

- ▶ We closely follow a technique introduced in an earlier version of the paper of Nazarov, Sodin & Volberg, but several new ideas are required to carry out the steps.
- ▶ For $L > 0$ and $x \in \mathbb{R}^d$ denote by $Q(x, L)$ the box $x + [-L, L]^d$.

Sketch of the proof

- ▶ We closely follow a technique introduced in an earlier version of the paper of Nazarov, Sodin & Volberg, but several new ideas are required to carry out the steps.
- ▶ For $L > 0$ and $x \in \mathbb{R}^d$ denote by $Q(x, L)$ the box $x + [-L, L]^d$.
- ▶ Let E_R denote the event that there exists an integral curve $\Gamma(t)$ connecting $\partial Q(0, R)$ and $\partial Q(0, 2R)$.

Sketch of the proof

- ▶ We closely follow a technique introduced in an earlier version of the paper of Nazarov, Sodin & Volberg, but several new ideas are required to carry out the steps.
- ▶ For $L > 0$ and $x \in \mathbb{R}^d$ denote by $Q(x, L)$ the box $x + [-L, L]^d$.
- ▶ Let E_R denote the event that there exists an integral curve $\Gamma(t)$ connecting $\partial Q(0, R)$ and $\partial Q(0, 2R)$.
- ▶ Easy to see: If X is the diameter of the basin containing 0, then for all $R > 0$,

$$\mathbb{P}(X \geq cR) \leq \mathbb{P}(E_R)$$

for some constant c depending only on the dimension.

- ▶ Introduce a gravitational potential energy function $U(x)$ whose differences $U(x) - U(y)$ are the line integrals of the gravitational force.

- ▶ Introduce a gravitational potential energy function $U(x)$ whose differences $U(x) - U(y)$ are the line integrals of the gravitational force.
- ▶ Most important observation: Since $F(x) = -\nabla U(x)$, therefore

$$\begin{aligned}\frac{d}{dt}U(\Gamma(t)) &= \langle \dot{\Gamma}(t), \nabla U(\Gamma(t)) \rangle \\ &= \langle F(\Gamma(t)), -F(\Gamma(t)) \rangle = -|F(\Gamma(t))|^2.\end{aligned}$$

In particular, the potential always decreases along a gradient flow.

Sketch contd.

- ▶ We bound $\mathbb{P}(E_R)$ in terms of discrete events, by dividing space into a grid S of cubes of side length $r \approx (\log R)^{2/d}$.

Sketch contd.

- ▶ We bound $\mathbb{P}(E_R)$ in terms of discrete events, by dividing space into a grid S of cubes of side length $r \approx (\log R)^{2/d}$.
- ▶ We fix $B = R^\alpha$ for some appropriate $\alpha < 1$ and say that a cube is **bad** if $|U(x) - U(y)| \leq Br/R$ for all x, y in the cube, or $U(x) < -B$ for some x in the cube.


Sketch contd.

- ▶ We bound $\mathbb{P}(E_R)$ in terms of discrete events, by dividing space into a grid S of cubes of side length $r \approx (\log R)^{2/d}$.
- ▶ We fix $B = R^\alpha$ for some appropriate $\alpha < 1$ and say that a cube is **bad** if $|U(x) - U(y)| \leq Br/R$ for all x, y in the cube, or $U(x) < -B$ for some x in the cube. (Essentially, a cube is bad if either the gravitational force or the potential is unusually small inside the cube.)

Sketch contd.

- ▶ We bound $\mathbb{P}(E_R)$ in terms of discrete events, by dividing space into a grid S of cubes of side length $r \approx (\log R)^{2/d}$.
- ▶ We fix $B = R^\alpha$ for some appropriate $\alpha < 1$ and say that a cube is **bad** if $|U(x) - U(y)| \leq Br/R$ for all x, y in the cube, or $U(x) < -B$ for some x in the cube. (Essentially, a cube is bad if either the gravitational force or the potential is unusually small inside the cube.)
- ▶ If $U(x) \leq B$ for all $x \in Q(0, 2R)$ (a high probability event for a right choice of α), then E_R happens only if there is a sequence of R/r *connected* bad cubes.

Schematic illustration



Here T_1 is the time at which the curve Γ enters the phase $U(x) < -B$.

Final steps

- ▶ Step 1: Show that the probability of a cube being bad is small, by further subdividing into smaller cubes and applying Markov's inequality to get a crude bound.

Final steps

- ▶ Step 1: Show that the probability of a cube being bad is small, by further subdividing into smaller cubes and applying Markov's inequality to get a crude bound.
- ▶ Step 2: Get an exponentially small bound on the joint probability of a collection of well-separated cubes being bad (**the percolation step**).

Final steps

- ▶ Step 1: Show that the probability of a cube being bad is small, by further subdividing into smaller cubes and applying Markov's inequality to get a crude bound.
- ▶ Step 2: Get an exponentially small bound on the joint probability of a collection of well-separated cubes being bad (**the percolation step**).
- ▶ Step 3: Bound $\mathbb{P}(E_R)$ by summing over all connected sequences of cubes.

Final steps

- ▶ Step 1: Show that the probability of a cube being bad is small, by further subdividing into smaller cubes and applying Markov's inequality to get a crude bound.
- ▶ Step 2: Get an exponentially small bound on the joint probability of a collection of well-separated cubes being bad (**the percolation step**).
- ▶ Step 3: Bound $\mathbb{P}(E_R)$ by summing over all connected sequences of cubes.
- ▶ The percolation step turns out to be rather difficult, mainly due to the heavy-tailed nature of the force (unlike the Nazarov-Sodin-Volberg scenario, where the dependence decreases exponentially with distance).

Sketch of the percolation step

The percolation step states, in essence, that the probability of the potential difference being simultaneously small across a large number of well-separated squares is exponentially small in the number of squares. The key component in the proof is the following bound on the joint density of the forces at multiple points.

Sketch of the percolation step

The percolation step states, in essence, that the probability of the potential difference being simultaneously small across a large number of well-separated squares is exponentially small in the number of squares. The key component in the proof is the following bound on the joint density of the forces at multiple points.

Lemma

Suppose we have $x_1, \dots, x_N \in \mathbb{R}^d$ with $|x_i - x_j| > S$ for every $i \neq j$. Fix $\lambda > 0$, and let

$$E = \left\{ \text{There is at least one star in } B(x_i, \lambda) \text{ for every } 1 \leq i \leq N \right\}.$$

Let \mathcal{M} the σ -algebra generated by the stars in $(\cup_{i=1}^N B(x_i, S))^c$. Then there exist constants $c_0, C_1 > 0$ such that if $\lambda < c_0 S (\log N)^{-1/d}$, then conditioned on the event E and on the σ -algebra \mathcal{M} , almost surely the joint density of $(F(x_i))_{1 \leq i \leq N}$ exists and is bounded from above by $(C_1 \lambda^{d^2-d})^N$.

- ▶ Proof of the lemma involves actual computation and bounding of the inverse Jacobian.

- ▶ Proof of the lemma involves actual computation and bounding of the inverse Jacobian.
- ▶ The bound is probably suboptimal, but suffices for our purposes.

- ▶ Proof of the lemma involves actual computation and bounding of the inverse Jacobian.
- ▶ The bound is probably suboptimal, but suffices for our purposes.
- ▶ Even after this lemma, many other technical hurdles need to be overcome.

- ▶ Proof of the lemma involves actual computation and bounding of the inverse Jacobian.
- ▶ The bound is probably suboptimal, but suffices for our purposes.
- ▶ Even after this lemma, many other technical hurdles need to be overcome.
- ▶ Dimensions 3 and 4 are harder, mainly because the invariant potential is no longer well-defined, and the technicalities are also more difficult.