

Instantons  
§  
Lagrangians  
in

Manifolds w/ V.C.P.  
[Vector Cross Product].

Naichung Conan Leung.

Joint work with JaeHyuk Lee.

- Vector Cross Product (V.C.P.)

- Kähler / Symplectic Manifold

$$\partial(\text{Holomorphic Curve}) \subset \text{Lagrangian Submanifold.}$$

- $G_2$  - manifold.

$$\partial(\text{Associative submanifold}) \subset \text{Coassociative submanifold.}$$

- $\mathbb{C}$  - V. C. P.

- Calabi - Yau Manifold.

$$\partial(\text{SLag}_\theta) \subset \begin{array}{l} \text{Complex Hypersurface} \\ \text{and} \\ \text{SLag}_{\theta + \pi/2}. \end{array}$$

- Hyperkähler Manifold.

$$\partial(\text{Holomorphic Curve}) \subset \text{Complex Lagrangian submanifold.}$$

# § Symplectic Geometry

## [Review].

$$(M^{2n}, \omega) \quad \omega \in \Omega^2(M).$$

$$\begin{cases} d\omega = 0 \\ \omega > 0. \end{cases}$$

Example.

$$M = \mathbb{C}^n$$

$$\omega = dx^1 \wedge dy^1 + \dots + dx^n \wedge dy^n$$

metric

$$g = (dx^1)^2 + (dy^1)^2 + \dots + (dx^n)^2 + (dy^n)^2$$

complex str.

$$J \left( \frac{\partial}{\partial x^i} \right) = \frac{\partial}{\partial y^i} \quad \forall i.$$

$$\omega \in \Omega^2(M)$$

complex structure.  
(1-fold v.c.p.)  
 $J: TM \rightarrow TM, J^2 = -$



via metric

$$\underline{\omega(u, v) = g(Ju, v)}$$

$(M, \omega, J, g)$

Almost Kähler mfd.

$$(M^{2n}, g, \omega, J)$$

Definition:  $A^2 \subset M$

Holomorphic Curve / Instanton

if  $A$  preserved by  $J$ .

[ $\Longleftrightarrow$  Wrtinger  $A$  calibrated by  $\omega$ , i.e.  $\omega|_A = \lambda A$ ]

[ $\implies$  Absolute minimum area.]

$$\text{Area}(A) = \int_A [\omega] \quad (\text{topological})$$

Remark: Boundary value problem

$$\partial A \subset C \subset M$$

$$\omega|_C = 0.$$

$$(M^{2n}, g, \omega, J)$$

Definition:  $C^n \subset M^{2n}$

Lagrangian submanifold if

$$\omega|_C = 0$$

$$\dim C = \frac{1}{2} \dim M.$$

Remark:

$$\# \{ \text{instantons } A^2 \subset M \}$$

(i)  $\partial A = \emptyset \rightsquigarrow$  Gromov Invariants - Witten

(ii)  $\partial A \subset C \rightsquigarrow$  Fukaya-Floer Category.

## §. Vector Cross Product.

Example:  $(\mathbb{R}^3, \times)$ .Example:  $(M^{2n}, J)$ .Definition:  $(M^m, g)$  Riemannian manifold.  
[Gray].

$$\chi: \Lambda^r T_M \rightarrow T_M \quad \underline{r\text{-fold VCP}}$$

if. (i)  $\chi(v_1, \dots, v_r) \perp v_i$

(ii)  $v_1, \dots, v_r$  : orthonormal  
 $\Rightarrow |\chi(v_1, \dots, v_r)| = 1$

(i)  $\Leftrightarrow$  (i)'  $\varphi(v_1, \dots, v_r, v_{r+1}) \triangleq \langle \chi(v_1, \dots, v_r), v_{r+1} \rangle$

then  $\varphi \in \Omega^{r+1}(M)$ .

(iii)  $d\varphi = 0$ .

[ (iii)  $\nabla\varphi = 0$  Integrable VCP ]

$(M^m, g)$  w/  $r$ -fold VCP.

$$\varphi \in \Omega^{r+1}(M) \quad \chi: \wedge^r T_M \rightarrow T_M$$

$$\underline{\varphi(v_1, \dots, v_r, v_{r+1}) = \langle \chi(v_1, \dots, v_r), v_{r+1} \rangle.}$$

Def<sup>n</sup>.  $A^{r+1} \subset M^m$  instanton

if  $A$  preserved by  $\chi$

[ Lemma.  $\iff$   $A$  calibrated by  $\varphi$  ]

Def<sup>n</sup>.  $C^k \subset M^m$  Lagrangian

if  $\varphi|_C = 0$

$$\dim C = k = \frac{n+r-1}{2}.$$

Remark:  $\dim C > \frac{n+r-1}{2} \Rightarrow \varphi|_C \neq 0.$



# § (Unparametrized) Loop Space Interpretation.

$$(M, g) \quad \varphi \in \Omega^{r+1}(M).$$


Fix ANY  $\Sigma^{r-1}$

$$\mathcal{L}_{\Sigma} M \triangleq \text{Map}(\Sigma, M)_{\text{embed}} / \text{Diff}^+(\Sigma).$$

Transgression  $\rightarrow$

$$\omega_{\mathcal{L}_{\Sigma} M} = \int_{\Sigma} \varphi \in \Omega^2(\mathcal{L}_{\Sigma} M).$$

That is,





$$\begin{aligned} & \omega_{\mathcal{L}_{\Sigma} M}(u, v) \\ &= \int_{\Sigma} \iota_{u \wedge v} \varphi \end{aligned}$$

Theorem.  $(M^n, g) \quad \varphi \in \Omega^{r+1}(M)$   
 $\rightsquigarrow \quad \omega_{\mathcal{L}_\Sigma M} \in \Omega^2(\mathcal{L}_\Sigma M).$

(1)  $\varphi$ :  $r$ -fold VCP on  $M$   $\iff$   $\omega_{\mathcal{L}_\Sigma M}$ : 1-fold VCP on  $\mathcal{L}_\Sigma M$ .  
 (i.e. Symplectic).

(2)  $C \subset M$   $\iff$   $\mathcal{L}_\Sigma C \subset \mathcal{L}_\Sigma M$   
 $\varphi$ -Lagrangian Lagrangian  
 $(\dim C = (n+r-1)/2).$

(3)  $D^2 \times \Sigma \subset M$   $\iff$   $D^2 \subset \mathcal{L}_\Sigma M$   
 $\parallel$   
 $A^{r+1}$   
 instanton instanton  
 (i.e. homomorphic curve)



# § Examples / Classification.

1° Kähler / Symplectic Manifolds.  
( $r = 1$ ).

$$(M^{2m}, \chi, \varphi) = (M, \underset{\substack{\uparrow \\ \text{cpx.} \\ \text{str.}}}{J}, \underset{\substack{\uparrow \\ \text{sympl.} \\ \text{form.}}}{\omega})$$

- Instanton (usual def<sup>n</sup>.)
- Lagrangian

2° Volume form ( $r = n - 1$ )

$$(M^n, g)$$

$$\begin{aligned} \varphi &= \nu_M \in \Omega^n(M) \\ &= \sqrt{\det(g_{ij})} dx^1 \wedge \dots \wedge dx^n \end{aligned}$$

• Instantons  $A^n \underset{\text{open}}{\subseteq} M^n$  (domain in  $M$ )

• Lagrangians  $C^{n-1} \subset M^n$

||  
Hypersurfaces

3°  $G_2$ -manifold ( $r=2$ ).

$$(M^7, g) \quad \varphi =: \Omega \in \Omega^3(M^7)$$

$$\left[ \begin{array}{l} \text{e.g. } M^7 = X^6 \times S^1 \\ \quad \uparrow \\ \quad \text{Calabi-Yau 3-fold.} \\ \Omega_M = \text{Re } \Omega_X + \omega_X \wedge d\theta \end{array} \right.$$

When  $M^7 = \text{Im } \mathbb{O}$

$$\chi(u, v) =: u \times v$$


$$= \text{Im}(u \cdot v)$$

$(M^7, g, \Omega) : G_2\text{-mfd.}$

$\Omega(u, v, w) = g(u \times v, w).$

• Instanton  $A^3 \subset M$  preserved by  $\chi$   
 || (calib. by  $\Omega$ )  
 Associative

• Lagrangian  $C^4 \subset M$   $\Omega|_C = 0$   
 || (calib. by  $*\Omega$ )  
 Coassociative.



Remark: Count  $A^3$ .  
 When  $C_1 \sim C_2$ . (L. - X.W. Wang)

$\# A^3 \approx \# \text{holo. curves in } C_1$  (4 bubbles).

Taubes  
 $\approx$  Seiberg-Witten Inv. of  $C_1$ .

# 4. Spin(7)-manifold $(r=3)$ $M^8$ .

$$\left[ \begin{array}{l} \text{eg. } M^8 = Z^7 \times S^1 \\ \quad \quad \quad \uparrow \\ \quad \quad \quad \mathbb{G}_2\text{-mfd.} \\ \varphi_M = \Omega_Z \wedge d\theta + * \Omega_Z. \end{array} \right.$$

Example:  $M = \mathbb{D}$

$$u \times v \times w = \mathcal{X}(u, v, w)$$

$$= \frac{1}{2} [u(\bar{\nabla} w) - w(\bar{\nabla} u)]$$

- Instanton  $A^4 \subset M^8$  preserved by  $\times$   
 $\parallel$  (calib. by  $\varphi$ )  
 Cayley

- Proposition:  $\nexists$   $\varphi$ -Lagrangian  
 in any Spin(7)-manifold.

Remark: No other VCP.—  
 classification by Brown-Gray.

In fact, we can also allow

$$\underline{r = 0} \text{ . i.e.}$$

$$(M^n, g) \quad \varphi \in \Omega^1(M)$$

$$\begin{cases} d\varphi = 0 \\ |\varphi| = 1. \end{cases}$$

Eg.

$$f: M \longrightarrow S^1 \quad \text{Riemannian submersion.}$$

$$\varphi = f^*(d\theta).$$

- Instanton  $A^1 \subset M$   
 $\parallel$   
 Gradient Flow Line.
- Lagrangian ?


$$f: M^n \rightarrow S^1$$

$\iff$

$$M = X \times [0, 1] / \sim$$

$$h: X \xrightarrow{\text{isometry}} X$$

(Mapping Cylinder).



• Lagrangian.

$$C^n \subset X^{2n} \times \{t\} \quad \exists t.$$




[  $r=0$  continue .... ]

$M^{2m+1}$

$\varphi \in \Omega^1(M)$

$f \downarrow$   
 $S^1$

$\parallel$   
 $f^*(d\theta)$



# Instantons bounding  $C_1 \cup C_2$

$$= \sum_{k=-\infty}^{\infty} \left( \# C_1 \cap h^k(C_2) \right) \pm^k.$$

Remark:  $(M^n, g), \varphi \in \Omega^{r+1}(M)$   
 VCP BUT  $d\varphi \neq 0$

Assume

(\*)  $| \iota_{u_1 \wedge \dots \wedge u_{r+1}}(d\varphi) |^2 = 1 - | \varphi(u_1, \dots, u_{r+1}) |^2$

∗ orthormal  $u_1, \dots, u_{r+1}$ .

- (\*)  $\Rightarrow \mathbb{R}_+ \times M$  (cone).  
 (r+1)-fold V.C.P.

$\tilde{\varphi} = dt \wedge \varphi + t \wedge d\varphi, \quad d\tilde{\varphi} = 0$

$\tilde{g} = t^2 g_M + dt^2$ .

- | $r$   | $M$ w/ (*)          |
|-------|---------------------|
| 0     | Contact             |
| 1     | Nearly C.Y. 3-fold. |
| 2     | Nearly $G_2$ -mfd.  |
| $n-1$ | Volume form.        |

# Remark (Continue)

$(M^{2m+1}, g)$      $\varphi \in \Omega^1(M)$      $\boxed{\gamma=0}$   
 $d\varphi$  satisfies  $(*)$

$\Rightarrow M$  : Contact.

• Instanton     $A' \subset M$

• Lagrangian     $C^m \subset M^{2m+1}$   
   ||  
 Legendrian

## § Complex V.C.P.

Definition:  $(M^{2n}, g, J)$ . Kähler.

$$\varphi \in \Omega^{r+1,0}(M), \quad d\varphi = 0$$

satisfying

$$|\int u_1 \wedge \dots \wedge u_r \varphi| = 1$$

for any orthonormal  $u_1, \dots, u_r \in T^{1,0}M$ .

is called a Complex Vector Cross Product.

## Classification Theorem (Lee-L.)

$$(M^{2n}, g, J), \quad \varphi \in \Omega^{r+1,0}(M)$$

$r$ -fold  $\mathbb{C}$ -V.C.P.

$\Rightarrow$  (1). Calabi-Yau. ( $r = n - 1$ ).

$\varphi =$  holomorphic Volume form

(2) Hyperkähler ( $r = 1$ ).

$\varphi =$  holomorphic Symplectic form.

Remark:  $\nexists$   $\mathbb{C}$ -analog. of  $G_2$  or  $Spin(7)$   
type V.C.P.

Remark:  $d\varphi = 0 \iff \nabla\varphi = 0$   
for  $\mathbb{C}$ -V.C.P.

$$(M^{2n}, J, g), \varphi \in \Omega^{\overline{r+1,0}}(M) \\ \subset VCP.$$

Def<sup>n</sup> (i)  $A^{2\overline{r+1}} \subset M$  Instanton

if Calibrated by  $\operatorname{Re}(e^{i\theta}\varphi)$   
for some 'phase'  $\theta$ .

(ii)  $C^{2k} \subset M^{2n}$  Lagrangian  
 $k = (n+r-1)/2$  (of ex. type).

if  $\varphi|_C = 0$ .

Remark:  $C \subset M$  Lagr.  $\Rightarrow$  complex submfd

Def. Lagrangian of Real Type

$$(M^{2n}, g, J)$$

$$\downarrow$$

$$\underline{\omega \in \Omega^2(M)}$$

$$\underline{\varphi \in \Omega^{r+1,0}(M)}$$

$$C^n \subset M^{2n}$$

$$(i) \omega|_C = 0$$

$$(ii) \operatorname{Re}(e^{i\theta}\varphi)|_C = 0$$

real type

$$C^{2k} \subset M^{2n}$$

$$\varphi|_C = 0$$

complex type

Remark: Both are good boundary value for Instantons.  
(Calib. by  $\operatorname{Re}(e^{i\theta}\varphi)$ ).

48. Example [ $r = 1$ ]: Hyperkähler.

$$(M^{4m}, g, J, \omega = \omega_J)$$

$$\varphi \in \Omega^{2,0}(M)$$

$$\parallel \\ \omega_I + i\omega_K.$$

- Instanton.  $A^2 \subset M$  calib. by  $\text{Re}(e^{i\theta}\varphi)$ .  
 $\parallel$   
 $J_\theta$ -holomorphic curve.

$$J_\theta = \cos\theta I + \sin\theta K.$$

- Lagr. of Cx. Type.  $C^{2n} \subset M^{4n}$   $\varphi|_C = 0$   
 $\parallel$   
J-Complex Lagr.

- Lagr. of Real Type.  $C^{2n} \subset M^{4n}$   $\omega|_C = 0$   
 $\parallel$   $\text{Re}(e^{i\theta}\varphi)|_C = 0$   
 $J_{\theta+\pi/2}$ -complex Lagr.



# Example [ $r = n - 1$ ] Calabi-Yau

$$(M^{2m}, g, J, \omega)$$

$$\varphi \in \Omega^{n,0}(M)$$

holo. vol. form.

- Instanton  $A^m \subset M^{2m}$  calib. by  $\operatorname{Re}(e^{i\theta}\varphi)$   
 $\parallel$


SLag w/ phase =  $\theta$ .

- Lagr. of  $\mathbb{C}$ -type  $C^{2m-2} \subset M^{2m}$   $\varphi|_C = 0$   
 $\parallel$   
 Complex Hypersurface.

- Lagr. of  $\mathbb{R}$ -type  $C^m \subset M^{2m}$   $\omega|_C = 0$   
 $\operatorname{Re}(e^{i\theta}\varphi)|_C = 0$   
 $\parallel$

SLag. w/ phase =  $\theta + \frac{\pi}{2}$ .

Remark:  $M^{2m} : C.Y.$



$SLag_{\pi/2}$

(Lagr.,  $\mathbb{R}$ -type)

Cx. Hypersurface.

(Lagr.  $\mathbb{C}$ -type).

$SLag_0$  with boundary lying on  $SLag_{\pi/2}$  (Dirichlet) or Cx. Hypersurface (Neuman)

— Schoen's school.  
(A. Butscher, W.Y. Qiu).


54.  
A

Remark: Motivation of Lagr.  
of  $\mathbb{R}$  vs  $\mathbb{C}$  type.  
(Complexification).

$$G_{\mathbb{C}}^{2n-2} = \{f(z)=0\} \subset \mathbb{C}^n$$

cx. type.      real type.

$$\mathbb{C}^n_{\mathbb{R}} = N_{\mathbb{C}/\mathbb{R}}^* \mathbb{R}^n$$



$(\mathbb{R}^n, g)$   
 $\omega$ : vol. form  
[ $r=n-1$ ]  
VCP.

# § Loop Space Interpretations of $\mathbb{C} \cdot \text{VCP}$ .

$$\left[ \begin{array}{l}
 \text{Recall: } [r = n - 1] \\
 (M^n, g), \quad \mu \in \Omega^n(M) \text{ (} n\text{-fold)} \\
 \Downarrow \\
 \mathcal{L}_{\Sigma^{n-2}} M, \quad \omega_{\mathcal{L}_{\Sigma} M} \in \Omega^2(\mathcal{L}_{\Sigma} M) \\
 \parallel \\
 \frac{\text{Map}(\Sigma^{n-2}, M^n)}{\text{Diff}(\Sigma)} \quad \text{1-fold (symplectic)}
 \end{array} \right.$$

262.  
A

## PLAN

$M^{2n}$  : Calabi-Yau.  $(n-1)$ -fold  $\mathbb{C}VCP$   
 $(M, g, J, \omega)$ ,  $\varphi \in \Omega^{n,0}(M)$



$$\frac{\text{Map}(\Sigma^{n-2}, M)}{\text{Diff}(\Sigma)}$$

: Hyperkähler

Symplectic Quotient ??

- For  $\omega/M$  induces a symplectic str.

$\omega_{\text{Map}}$  on  $\text{Map}(\Sigma, M)$ . Need

**FIX.**  $\nu_{\Sigma} \in \Omega^{n-2}(\Sigma)$ . i.e.

$$\omega_{\text{Map}}(u, v) = \int_{\Sigma} \omega(u, v) \nu_{\Sigma}$$

- Problem:  $\omega_{\text{Map}}$  is preserved only by  $\text{Diff}(\Sigma, \nu_{\Sigma})$ , not  $\text{Diff}(\Sigma)$ .

## Moment Map (Donaldson, Hitchin)

$$\text{Diff}(\Sigma, \omega_\Sigma) \xrightarrow{\quad} \text{Map}(\Sigma, M) \xrightarrow{\mu} \Omega^1(\Sigma) / d\Omega^0(\Sigma)$$

$$\mu(\Sigma \xrightarrow{f} M) = \alpha$$

$$\text{where } d\alpha = f^* \omega.$$

$$\frac{\mu^{-1}(0)}{\text{Diff}(\Sigma, \omega)}$$

: Symplectic  
(NOT Hyperkähler)

$$\frac{\text{Map}(\Sigma, M)}{\text{Diff}(\Sigma, \omega)}$$

Want • // Diff( $\Sigma$ )

56.  
A

AIM:  $(M^{2n}, \omega \in \Omega^{1,1}, \varphi \in \Omega^{n,0}) : \text{C.Y.}$   
 Construct Hyperkähler " $\text{Map}(\Sigma^{n-2}, M)$ "  
 $\text{Diff}(\Sigma)$

$$\tilde{\varphi} := \tilde{\omega}_I - i\tilde{\omega}_K \stackrel{\text{def}}{=} \int_{\Sigma} \varphi \quad (2,0)\text{-form on } \text{Map}(\Sigma, M).$$

$\mapsto I, K : T(\text{Map}) \ni$

BUT  $\begin{cases} I^2 \neq -1 \neq K^2 \\ \tilde{\omega}_I, \tilde{\omega}_K : \text{degenerate.} \end{cases}$

$\tilde{\omega}_I$ -Sympl. Reduction of  $\mu^{-1}(0)$   
 $\parallel$   
 $\{\Sigma \hookrightarrow M \text{ isotropic}\}$

$\tilde{\omega}_K$ -Sympl. Reduction of  $\mu^{-1}(0)$

is Hyperkähler !

$(\tilde{\omega}_I, \omega_{\text{Map}}, \tilde{\omega}_K)$   
 $\parallel$   
 $\omega_J$

$\text{Map}(\Sigma, M) // \text{Diff}(\Sigma).$

068  
A  $M^{2n}$  Calabi-Yau.  $\rightsquigarrow$   $\tilde{\mathcal{L}}_{\Sigma} M$  Hyperkähler  
 "  $\text{Map}(\Sigma^n; M) // \text{Diff}(\Sigma)$  "

$\omega \in \Omega^2(M) \rightsquigarrow \tilde{\omega}_J \in \Omega^2(\tilde{\mathcal{L}}_{\Sigma} M)$

$\varphi \in \Omega^{n,0}(M)$   $\rightsquigarrow$   $\tilde{\varphi} \in \Omega^{2,0}(\tilde{\mathcal{L}}_{\Sigma} M)$   
 holo. volume form  $\tilde{\omega}_J - i \tilde{\omega}_K$  holo. sympl. form

$A^n \subset M$   $\longleftrightarrow$   $D^2 \subset \tilde{\mathcal{L}}_{\Sigma} M$   
 $D^2 \times \Sigma$  instanton (i.e.  $\text{SLag}_{\theta + \pi/2}$ ) instanton (i.e. holo. curve).

$C^{2n-2} \subset M$   $\longleftrightarrow$   $\tilde{\mathcal{L}}_{\Sigma} C \subset \tilde{\mathcal{L}}_{\Sigma} M$   
 Lagr.  $\mathbb{C}$ -type. J-complex Lagr.  
 (i.e.  $C_x$  hypersurface).

$C^n \subset M^{2n}$   $\longleftrightarrow$   $\tilde{\mathcal{L}}_{\Sigma} C \subset \tilde{\mathcal{L}}_{\Sigma} M$   
 Lagr.  $\mathbb{R}$ -type  $\text{Map}(\Sigma, C) // \text{Diff}(\Sigma)$  automatic  
 (i.e.  $\text{SLag}_{\theta}$ ).  $\text{J}\theta$ -complex Lagr.



270. Remark: MCF (Mean Curv. Flow) on manifolds w/  $\mathbb{C}$ -VCP (C.Y. & H.K.).

Calabi-Yau case:  $C^n \subseteq M^{2n}$ : CY.

$$\omega|_C = 0$$

(i.e.  $\mathbb{R}$ -Lagr.)

$$\iff T_x C \subset T_x M \text{ calibrated by } \operatorname{Re}(e^{i\theta(x)} \varphi(x))$$

for some  $\theta(x): C \rightarrow S^1$ .

$$\Rightarrow \int_H \omega = d\theta$$

$$\Rightarrow \left\{ \begin{array}{l} \text{MCF preserve Lagr.} \\ \text{Gr. pt.: } \theta \equiv \text{const (sLagr.)} \end{array} \right.$$

072  
A

$(M^{4n}, \omega_J, \Omega_J = \omega_I + i\omega_K)$   
Hyperkähler.

If  $C^{2n} \subset M^{4n}$  st.  $\forall x \in C$

$T_x C \subset T_x M$  Calib. by  $\Omega_{J\theta(x)}$

$\exists \theta(x) \in S^2_{\text{twistor}}$ .

then  $C$  is called a  $\mathbb{C}$ -Lagrangian

Theorem (L. and Tom Wan).

•  $L_{\bar{H}} \Omega_J = \partial_J \theta$

$J = J_{\theta(x)}$ .

• MCF preserves  $\mathbb{C}$ -Lagr.

• Min.  $\mathbb{C}$ -Lagr.  $\Rightarrow \theta: C \rightarrow S^2$   
anti-holomorphic.

•  $\theta(x) \equiv \text{const.} \Leftrightarrow$  Complex Lagrangian (calibrated)  
(i.e. Min.  $\mathbb{C}$ -Lagr. +  $[\theta(x)] = 0 \in [C, S^2]$ ).

# Comparisons:

## (1) Vector Cross Product

$r =$	$n-1$	$n=1$	$2$	$3$
VCP	Oriented	Kähler	$G_2$ -mfd	Spin7-mfd
$\mathbb{C}$ -VCP	Calabi-Yau	Hyperkähler		

## (2) Geometry / Normed Algebras $\mathbb{A}$ .

$\mathbb{A} =$	$\mathbb{R}$	$\mathbb{C}$	$\mathbb{H}$	$\mathbb{O}$
$\mathbb{A}$ -mfd	Manifold	Kähler	Quaternionic Kähler	Spin7-mfd
Special $\mathbb{A}$ -mfd	Oriented	Calabi-Yau	Hyperkähler	$G_2$ -mfd